

Silencer® Select Human Protease siRNA Library V4 Product Information Sheet

Product:	Silencer® Select Protease siRNA Library V4
Catalog # (P/N):	4397917
Lot #:	XXXXXXXX
Amount:	0.25 nmol each siRNA
Content:	1482 unique siRNAs targeting each of 494 human protease genes* Total of 18 96-well plates (plates are Axygen #PCR96FS; www.axxygen.com) 15 plates with 88 siRNAs each 3 plates with 54 siRNAs each *A few siRNAs have more than one gene target, due to gene families with highly homologous members or predicted genes with high homology to verified genes. See the accompanying CD for annotation information and siRNA details.
Purity:	Standard
siRNA Format:	Annealed
Appearance:	Powder
Storage:	Store at or below –20°C. Do not store in a frost-free freezer. (Dried oligonucleotides are shipped at ambient temperature.)
Shelf Life:	1 year, when stored at or below –20°C.

USER INFORMATION

Product Description: This Silencer® Select siRNA Library is composed of Silencer Select Pre-designed and, where available, Validated siRNAs, targeting human protease genes identified using the PANTHER™ and Gene Ontology databases.

Silencer Select siRNAs are designed using a novel algorithm that was developed utilizing the latest in machine-learning methods. These next-generation siRNAs exhibit up to 100-fold higher silencing potency than siRNAs from other leading siRNA manufacturers. Off-target activity (assayed by microarray analysis) is blocked by up to 90% because Silencer Select siRNAs can be used at 5- to 20-fold lower concentrations, are bioinformatically screened using the latest knowledge about miRNA seed regions and toxic sequence motifs, and incorporate strategic chemical modifications. As a result, Silencer Select siRNAs provide unrivalled specificity and cleaner, more consistent phenotypic data.

Handling Instructions: RNA oligonucleotides are susceptible to degradation by exogenous ribonucleases introduced during handling. Wear gloves when handling this product. Use RNase-free reagents, tubes, and barrier pipette tips. Upon receipt, your siRNAs may be safely stored in a non-frost-free freezer at or below –20°C (dried oligonucleotides are shipped at ambient temperature).

Resuspension of siRNAs

Centrifuge each plate at low speed (maximum RCF 4,000 X g) to collect the contents at the bottom of the wells before removing the seal.

Important: Perform this process in a tissue culture hood.

1. Remove seal carefully.
2. Add nuclease-free, sterile water, using a multichannel pipette and sterile tips, to achieve the desired concentration.*
3. Gently pipet up and down 5 times to resuspend.
4. Centrifuge briefly to collect the liquid at the bottom of the wells, if necessary.
5. (Optional) Aliquot the siRNAs into one or more daughter plates, to limit the number of freeze-thaw cycles to which the siRNAs are subjected.
6. Place a new sterile seal (such as Axygen PCR-AS-200) on the plate before storing.
7. Store at –80°C until ready to use.

* An online calculator for suspension of dried oligonucleotides is available at www.ambion.com/techlib/append/oligo_dilution.html

Applications:**Transfecting Silencer Select siRNAs Into Mammalian Cells**

The efficiency with which mammalian cells are transfected with siRNA will vary according to cell type and the transfection agent used. This means that the optimal concentration used for transfections should be determined empirically. Since Silencer Select siRNAs exhibit superior silencing potency compared to other siRNAs, we suggest starting concentrations of 5- to 20-fold less than typically used for transfection of your experimental cell lines. We have found that Silencer Select siRNAs reduced mRNA levels >80% at final concentrations of 2–10 nM using lipid-mediated transfection in HeLa and U-2 OS human osteosarcoma cells.

General Transfection Starting Points for Mammalian Cells^a

Plate Format	384 wells	96 wells	24 wells	12 wells
Transfection Agent ^b	0.3 µL	0.3–1.0 µL	1–3 µL	2–4 µL
siRNA ^c	0.35–0.5 pmol	0.5 pmol	2.5 pmol	5 pmol
Cell Density ^d	4000–8000 cells/well	6000 cells/well	40,000 cells/well	80,000 cells/well
Final Volume per Well	70–100 µL	100 µL	0.5 mL	1.0 mL

^a Appropriate for lipid-mediated transfection and easily transfected cell lines such as HeLa.

^b Refer to the instructions provided with your transfection agent for the recommended volume.

^c The siRNA amounts indicated result in a final siRNA concentration of 5 nM. The amount of siRNA required for maximal gene silencing will vary among cell types. For a 96-well plate, and a 100 µL final transfection volume, 0.5 pmol of a 1 µM siRNA solution is 0.5 µL. Robotic pipettors may require volumes of 2–5 µL for accurate pipetting. To increase pipetting volumes and accuracy when preparing transfection complexes, we recommend first preparing a plate with a dilution of your stock siRNA.

^d Optimal cell density will vary among cell types, depending on cell size and growth characteristics. In general, 30–70% confluency is recommended.

Transfection Optimization

Optimizing transfection efficiency is crucial for maximizing gene silencing while minimizing cytotoxicity. Optimal transfection efficiencies are achieved by identifying an effective transfection agent for each cell type and by adjusting (in order of importance):

- Amount of transfection agent
- Amount of siRNA
- Cell density at the time of transfection
- Order of transfection (pre-plating cells or plating cells/transfecting in tandem)
- Length of exposure of cells to transfection agent/siRNA complexes

Most protocols recommend maintaining mammalian cells in the medium used for transfection; this avoids diluting or removing the siRNAs from the cells by adding medium or washing the cells with new medium too soon after transfection. We have found that cells typically exhibit greater viability when existing medium is replaced by fresh medium 24 hours after transfection. Replacing medium after 24 hours generally does not change the activity of the transfected siRNAs.

Once the conditions for maximal gene silencing are determined, they should be kept constant from experiment to experiment for a given cell type. Include controls in all plates for each experiment to ensure consistency.

For additional information about siRNA transfection, including transfection conditions for many cell types and optimization protocols, see the siRNA Delivery Resource at:
www.ambion.com/techlib/resources/delivery

RELATED PRODUCTS**Silencer® Select Pre-designed and Validated siRNAs**

Cat #Various (see www.ambion.com/geneassist)

An all-new class of modified siRNAs with unsurpassed efficacy, potency and specificity. Search the GeneAssist™ Atlas at www.ambion.com/geneassist to find guaranteed-to-silence siRNAs to your gene of interest.

Silencer® Select Control siRNAs

Cat #Various (see www.ambion.com/siRNA)

Validated, nontargeting siRNAs (Silencer Select Negative Controls) and proven-to-work positive control siRNAs (Silencer Select GAPDH siRNAs) with the Silencer Select modifications.

siPORT™ NeoFX™ Transfection Agent

Cat #AM4510 and AM4511

A versatile lipid-based agent for efficient and reproducible transfection of adherent cells while subculturing, without increased cytotoxicity.

TaqMan® Gene Expression Assays

See www.allgenes.com or www.ambion.com/geneassist

A comprehensive collection of over 700,000 probe and primer sets for quantitative gene expression analysis using real-time PCR. Search the GeneAssist™ Atlas at www.ambion.com/geneassist to find suggested

TaqMan Gene Expression Assays for the gene targeted by an siRNA of interest.

QUALITY CONTROL

Identity: The mass of a sample of each single-stranded RNA oligonucleotide is analyzed using MALDI-TOF mass spectrometry and compared to the calculated mass.

Annealing: A sample of the annealed siRNA is analyzed by nondenaturing gel electrophoresis.

OTHER INFORMATION

Material Safety Data Sheets: Material Safety Data Sheets (MSDSs) can be printed or downloaded from product-specific links on our website at the following address: www.ambion.com/techlib/msds. Alternatively, e-mail your request to MSDS_Inquiry_CCRM@appliedbiosystems.com. Specify the catalog or part number(s) of the product(s), and we will e-mail the associated MSDSs unless you specify a preference for fax delivery. For customers without access to the internet or fax, our technical service department can fulfill MSDS requests placed by telephone or postal mail. (Requests for postal delivery require 1–2 weeks for processing.)

Warranty and Liability: *For research use only. Not for use in diagnostic procedures.*

Applied Biosystems is committed to delivering superior product quality and performance, supported by industry-leading global service and technical support teams. Warranty information for the accompanying consumable product is available at www.ambion.com/info/warranty in "Limited Warranty for Consumables," which is subject to the exclusions, conditions, exceptions, and limitations set forth under the caption "EXCLUSIONS, CONDITIONS, EXCEPTIONS, AND LIMITATIONS" in the full warranty statement. Please contact Applied Biosystems if you have any questions about our warranties or would like information about post-warranty support.

Information in this document is subject to change without notice. Applied Biosystems assumes no responsibility for any errors that may appear in this document.

Applied Biosystems disclaims all warranties with respect to this document, expressed or implied, including but not limited to those of merchantability or fitness for a particular purpose. In no event shall Applied Biosystems be liable, whether in contract, tort, warranty, or under any statute or on any other basis for special, incidental, indirect, punitive, multiple or consequential damages in connection with or arising from this document, including but not limited to the use thereof.

Trademarks, Patents and Licensing:

Applied Biosystems, AB (Design), Ambion, and *Silencer* are registered trademarks and GeneAssist, *NeoFX*, PANTHER, and siPORT are trademarks of Applied Biosystems or its subsidiaries in the US and/or certain other countries. TaqMan is a registered trademark of Roche Molecular Systems, Inc. LNA is a registered trademark of Exiqon A/S, Vedbaek, Denmark. All other trademarks are the sole property of their respective owners.

NOTICE TO PURCHASER:

This product is licensed under European Patents 1144623, 1214945 and foreign equivalents from Alnylam Pharmaceuticals, Inc., Cambridge, USA and is provided only for use in academic and commercial research including in vitro and in vivo identification and validation of drug targets (but excluding the evaluation or characterization of this product as the potential basis for a siRNA-based drug) and not for any other commercial purposes. Information about licenses for commercial use (including discovery and development of siRNA-based drugs) is available from Alnylam Pharmaceuticals, Inc., 300 Third Street, Cambridge, MA 02142, USA.

Ambion siRNA products are manufactured under license from the Massachusetts Institute of Technology to U.S. Patent Nos. 7,056,704 and 7,078,196 and pending counterparts.

Purchase of this product gives the user rights under U.S. Patent No. 6,506,559, and its foreign equivalents, to use this product for life science research, not for use in humans or clinical diagnostics.

Locked Nucleic Acids are covered by patents and patent applications owned by Exiqon A/S. This product is for research use only and not for diagnostic or therapeutic use. The products may be used only for the buyer's internal research purposes and not for commercial use except for providing a service to research customers. The buyer may not resell products in their original or any modified form.

A license to perform the patented 5' Nuclease Process for research is obtained by the purchase of (i) both Licensed Probe and Authorized 5' Nuclease Core Kit, (ii) a Licensed 5' Nuclease Kit, or (iii) license rights from Applied Biosystems.

The TaqMan[®] Gene Expression Assays contain Licensed Probe. Use of the TaqMan Gene Expression Assays is covered by US patent claims and patent claims outside the US. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. The right to use this product in the 5' Nuclease Process under the applicable claims of US patents, and corresponding patent claims outside the United States, can be obtained through purchase of an Authorized 5' Nuclease Core Kit. Except under separate license rights available from Applied Biosystems, no right under any other patent claim, or to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, or to sublicense, repackage with other products, or resell in any form, is conveyed expressly, by implication, or by estoppel. This product is for research use only. Human in vitro diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained from the Director of Licensing, Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404, USA.

© 2008, 2010 Ambion, Inc. All rights reserved. 4398635B