CD71 (Transferrin Receptor) Monoclonal Antibody (OKT9 (OKT-9)), Functional Grade, eBioscience™

Size	50 µg
Species Reactivity	Human
Published Species	Human
Host/Isotype	Mouse / IgG1, kappa
Recommended Isotype Control	Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), Functional Grade, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	OKT9 (OKT-9)
Conjugate	Functional Grade
Form	Liquid
Concentration	1 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2
Contains	no preservative
Storage conditions	4° C
RRID	AB_469002

Applications	Tested Dilution	Publications
Immunocytochemistry (ICC/IF)	-	1 Publication
Flow Cytometry (Flow)	0.5 µg/test	14 Publications
Functional Assay (FN)	Assay-Dependent	-
Control (Ctrl)	Assay-Dependent	-

Product Specific Information

Description: The OKT9 monoclonal antibody reacts with human CD71, a 170-180 kDa type II transmembrane protein. CD71, the transferrin receptor, exists as a homodimer on the cell surface and is essential for cellular growth. CD71 is expressed by immature proliferating cells and at low levels on resting mature lymphocytes. Antigen or mitogen stimulation of T and B cells upregulates the expression of CD71.

Applications Reported: This OKT9 (OKT-9) antibody has been reported for use in flow cytometric analysis.

Applications Tested: This OKT9 (OKT-9) antibody has been tested by flow cytometric analysis of PHA-stimulated human peripheral blood mononuclear cells. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

1

Storage and handling: Use in a sterile environment.

Filtration: 0.2 µm post-manufacturing filtered.

Purity: Greater than 90%, as determined by SDS-PAGE.

Endotoxin Level: Less than 0.001 ng/µg antibody, as determined by LAL assay.

Aggregation: Less than 10%, as determined by HPLC.

□ 15 References

Immunocytochemistry (1)

Oncogene Terminal differentiation and loss of tumorigenicity of human cancers via pluripotency-based reprogramming. "Published figure using CD71 (Transferrin Receptor) monoclonal antibody (Product # 16-0719-81) in Immunofluorescence" Authors: Zhang X,Cruz FD,Terry M,Remotti F,Matushansky I	Species Not Applicable Dilution Not Cited Year 2013
Flow Cytometry (14)	
eLife	Species Not Applicable
Gene-centric functional dissection of human genetic variation uncovers	
regulators of hematopoiesis.	Dilution
"Published figure using CD71 (Transferrin Receptor) monoclonal antibody (Product # 16-0719-81) in Flow Cytometry"	Not Cited
Authors: Nandakumar SK,McFarland SK,Mateyka LM,Lareau CA,Ulirsch JC,Ludwig LS,Agarwal G,Engreitz JM, Przychodzen B,McConkey M,Cowley GS,Doench JG,Maciejewski JP,Ebert BL,Root DE,Sankaran VG	Year 2019
The Journal of biological chemistry	Species
Hormonal vitamin D up-regulates tissue-specific PD-L1 and PD-L2	Not Applicable
surface glycoprotein expression in humans but not mice.	Dilution
"Published figure using CD71 (Transferrin Receptor) monoclonal antibody (Product # 16-0719-81) in Flow Cytometry"	Not Cited
Authors: Dimitrov V,Bouttier M,Boukhaled G,Salehi-Tabar R,Avramescu RG,Memari B,Hasaj B,Lukacs GL,Krawczyk CM,White JH	Year 2017

View more Flow references on thermofisher.com

More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer's mereing liburative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTABILITY, INTERS FOR ANY PARTICLUOUTS DURING THE WARRANT PERFODE INTERVISE TO REPLAY. REPLACEMENT FOR NON-NORFINGEMENT, BUYER'S SCL. DEVIDENT OR REPLAY, DER CONCORTS DURING THE WARRANT PERFORD IN CLUDING WITHOUT LIMITATION, IMPLIED WARRANTES OF MERCHANTABILITY, FITNESS FOR ANY PARTICLUAR DURDES CR REFUNE FOR PRODUCTS SA STHE RESULT PORTONICTS ANY PARTICLUAR DURDES TO REPLAY. REPLACEMENT FOR REPLAY. DER CONCORTS DURING THE WARRANT PERFORD TO REPARE, REPLACEMENT FOR REPLAY. DER CONCORTS DURING THE WARRANT PERFORD USES AND THE RESULT OF (I) ACCLIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MSUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PONDUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. UNless otherwise expressly stated on the Product or in the documentation accompanying the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic use