# MHC Class II (I-A/I-E) Monoclonal Antibody (M5/114.15.2), eFluor 450, eBioscience™

Product Details	
Size	100 µg
Species Reactivity	Mouse
Published Species	Mouse
Host/Isotype	Rat / IgG2b, kappa
Recommended Isotype Control	Rat IgG2b kappa Isotype Control (eB149/10H5), eFluor 450, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	M5/114.15.2
Conjugate	eFluor® 450
Form	Liquid
Concentration	0.2 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_1272204

Applications	Tested Dilution	Publications
Immunohistochemistry (IHC)	-	16 Publications
Immunocytochemistry (ICC/IF)	-	3 Publications
Flow Cytometry (Flow)	0.125 µg/test	110 Publications
Immunoprecipitation (IP)	-	2 Publications

### Product Specific Information

Description: The M5/114.15.2 monoclonal antibody reacts with the mouse major histocompatibility complex class II, both I-A and I-E subregion-encoded glycoproteins (I-A b, I-A d, I-A q, I-E d, I-E k, not I-A f, I-A k, or I-A s). It detects a polymorphic determinant present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2 b, H-2 d, H-2 q, H-2 p, H-2 r and H-2 u but not from mice carrying the H-2 s or H-2 f haplotypes. The M5/114 mAb is reported to inhibit I-A-restricted T cell responses of the H-2 b, H-2 d, H-2 u but not H-2 d, H-2 u but not H-2 h, H-2 h,

Applications Reported: This M5/114.15.2 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This M5/114.15.2 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.125  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.


1

eFluor® 450 is an alternative to Pacific Blue®. eFluor® 450 emits at 445 nm and is excited with the Violet laser (405 nm). Please make sure that your instrument is capable of detecting this fluorochome.

Excitation: 405 nm; Emission: 445 nm; Laser: Violet Laser.

Filtration: 0.2 µm post-manufacturing filtered.

# Product Images For MHC Class II (I-A/I-E) Monoclonal Antibody (M5/114.15.2), eFluor 450, eBioscience™



## MHC Class II (I-A/I-E) Antibody (48-5321-82) in Flow

Staining of C57BL/6 splenocytes with Anti-Mouse CD3e PE (Product # 12-0031-82) and 0.06 µg of Rat IgG2b K Isotype Control eFluor® 450 (Product # 48-4031-82) (left) or 0.06 µg of Anti-Mouse MHC Class II (I-A/I-E) eFluor® 450 (right). Total viable cells were used for analysis.

View more figures on thermofisher.com

## □ 131 References

### Immunohistochemistry (16)

Frontiers in aging neuroscience	Species
Age Influences Microglial Activation After Cuprizone-Induced	Not Applicable
Demyelination.	Dilution
Published figure using MHC Class II (I-A/I-E) monoclonal antibody (Product # 48-5321-82) in Immunofluorescence"	Not Cited
Authors: Klein B,Mrowetz H,Barker CM,Lange S,Rivera FJ,Aigner L	<b>Year</b> 2020
Scientific reports	Species
•	Species Not Applicable
<sup>Scientific reports</sup> Intravitreal injection of adenosine A <sub>2A</sub> receptor antagonist reduces neuroinflammation, vascular leakage and cell death in the retina of diabetic mice.	

View more IHC references on thermofisher.com

# Immunocytochemistry (3)

<sup>Glia</sup> Exosomes derived from microglia exposed to elevated pressure amplify the neuroinflammatory response in retinal cells.	Species Not Applicable Dilution Not Cited
"Published figure using MHC Class II (I-A/I-E) monoclonal antibody (Product # 48-5321-82) in Immunocytochemistry"	
Authors: Aires ID,Ribeiro-Rodrigues T,Boia R,Catarino S,Girão H,Ambrósio AF,Santiago AR	<b>Year</b> 2020
Glia	Species
Functional antagonism of sphingosine-1-phosphate receptor 1 prevents	Not Applicable
cuprizone-induced demyelination.	Dilution
"Published figure using MHC Class II (I-A/I-E) monoclonal antibody (Product # 48-5321-82) in Immunocytochemistry"	Not Cited
Authors: Kim S,Bielawski J,Yang H,Kong Y,Zhou B,Li J	<b>Year</b> 2018

View more ICC/IF references on thermofisher.com

#### More applications with references on thermofisher.com

Flow (110) IP (2)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty periode herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer's mereing liburaris with the Product is subjected to normal, proper and intended uses. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer's mereing liburaris with the Product of the Goes not represent that any Product will conform to such model or sample. No OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTES OF MERCHANTABILITY, FITNESS FOR ANY PARTICLUAR PURPOSE, OR RON INFRINCEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONCINTS DURING THE WARRANT PERIOD IS LIMITED TO REPARE, REPLACEMENT OF OR REFUNE TO REPARE REPLACE OF OR REPUNE TO REPARE REPLACE OF OR BY BUYER, (III) USE OF THE NON-CONFORMING PRODUCTS SURIA BORDING TO RODUCTS DURING THE WARRANTES OF REPUNE TO REPARE AND REPLACE PARE DURING THE WARRANT REPLACEMENT OF FORCE MAJELINE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product; is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagn

3