

AxyPrep Multisource Genomic DNA Miniprep Kit

For the purification of genomic DNA from animal tissues, plant tissues, filamentous fungi, cultured cells, lymphocytes, bone marrow, bone, buccal swabs and yeast

Kit contents, storage and stability

Cat. No.	AP-MN- MS-GDNA-50	AP-MN- MS-GDNA-250
Kit size	50 preps	250 preps
Miniprep column	50	250
2 ml Microfuge tube	100	500
1.5 ml Microfuge tube	50	250
RNase A	60 µl	300 µl
Buffer C-L	9 ml	45 ml
Proteinase K	18 mg	90 mg
PK Buffer	1.2 ml	6 ml
Buffer P-D	25 ml	125 ml
Buffer W1	30 ml	145 ml
Buffer W2 concentrate	24 ml	120 ml
Eluent	12 ml	60 ml
Protocol manual	1	1

Except for the RNase A and Proteinase K, all buffers are stable for a period of at least 12 months from the date of receipt when stored under ambient conditions. Please avoid exposure to direct sunlight or extremes in temperature. After reconstitution, Proteinase K is stable for 2 months when stored at 4°C. To preserve RNase A activity, the RNase A is suspended in a solution containing a high concentration of ammonium sulfate. The lyophilized Proteinase K is resuspended in PK Buffer, which also contains ammonium sulfate. On occasion, a precipitate may form. If this occurs, resuspend by vortexing or pipetting before use. Enzymatic activity is unaffected by the formation of a precipitate.

RNase A: 50 mg/ml. Store at room temperature.

Buffer C-L: Lysis buffer. Store at room temperature.

Proteinase K: Lyophilized Proteinase K is stable for up to 6 months after delivery when stored at room temperature. After reconstitution with PK Buffer, Proteinase K is stable for 2 months when stored at 4°C. Storing the Proteinase K stock solution at room temperature for prolonged periods of time should be avoided.

PK Buffer: Used to resuspend Proteinase K. Store at room temperature.

Buffer P-D: Protein precipitation buffer. Store at room temperature.

Buffer W1: Wash buffer. Store at room temperature.

Buffer W2 concentrate: Desalting buffer. Before using the kit, add the amount of ethanol specified on the bottle label to the Buffer W2 concentrate. Either 100% or 95% denatured ethanol can be used. Store at room temperature.

Eluent: 2.5 mM Tris-HCl, pH 8.5. Store at room temperature.

Introduction

The AxyPrep Multisource Genomic DNA Miniprep Kit is designed to purify genomic DNA from a wide variety of different starting materials. This system employs a special lysis Buffer C-L and Proteinase K, to efficiently release genomic DNA from the biologic starting materials. Contaminating proteins, pigments, carbohydrates and lipids are then efficiently segregated from the genomic DNA by precipitation with Buffer P-D, coupled with the selective adsorption of the genomic DNA to a special AxyPrep column. After washing and desalting, the purified DNA is then eluted in either a Tris buffer or deionized water. Genomic DNA prepared by this method is approximately 30 Kb in length and is suitable for a variety of applications, such as PCR amplification, Southern blot analysis, etc. Each AxyPrep column will bind and purify up to 20 µg of genomic DNA.

For the purification of genomic DNA from whole blood, we recommend the AxyPrep Blood Genomic DNA Kits: Mini (#AP-MN-BL-GDNA-4/50/250); Midi (#AP-MD-BL-GDNA-2/10/25); Maxi (#AP-MX-BL-GDNA-2/10/25). For the purification of bacterial genomic DNA, please refer to the AxyPrep Bacterial Genomic DNA Miniprep Kit (#AP-MN-BT-GDNA-4/50/250).

Caution

Buffer C-L, Buffer P-D and Buffer W1 contain chemical irritants. When working with these buffers, always wear suitable protection such as safety glasses, laboratory coat and gloves. Be careful to avoid contact with eyes and skin. In the case of such contact, wash immediately with water. If necessary, seek medical assistance.

Equipment and consumables required

- Microcentrifuge capable of 12,000xg
- Mortar and pestle
- Heated water bath
- AxyVac Vacuum manifold (#AP-VM)
- Vacuum regulator
- Vacuum source (-25-30 inches Hg required)
- 100% or 95% (denatured) ethanol

Preparation before experiment

- 1) Reconstitution of Buffer W2. Before using the kit, add the amount of ethanol specified on the Buffer W2 label and mix well. Either 100% or 95% (denatured) ethanol can be used.
- 2) Adjust a water bath to 56°C.

- 3) Check Buffer C-L and Buffer P-D for precipitation before each use. If precipitation occurs, incubate at 37°C to dissolve the precipitate.
- 4) Pre-warming the Eluent to 65°C will often improve elution efficiency.
- 5) Resuspend Proteinase K in PK Buffer.
- 6) Prepare phosphate-buffered saline (PBS). Dissolve in 800 ml of distilled H₂O: 8 g of NaCl, 0.2 g of KCl, 1.44 g of Na₂HPO₄ and 0.24 g of KH₂PO₄. Adjust the pH to 7.4 with HCl. Add H₂O to 1 liter.

Protocols

- I. Purification of Genomic DNA from Animal Tissues
- II. Purification of Genomic DNA from Plant Tissues and Filamentous Fungi
- III. Purification of Genomic DNA from Cultured Cells, Lymphocytes, Bone Marrow and Bone
- IV. Purification of Genomic DNA from Buccal Swabs
- V. Purification of Genomic DNA from Yeast

Each type of starting materials has different requirements for the method(s) used to achieve efficient lysis and homogenization. Both vacuum and centrifuge protocols are provided for each starting material type.

I. Purification of Genomic DNA from Animal Tissues

[Vacuum protocol]

This procedure requires use of the AxyVac vacuum manifold or other manifold with complementary luer fittings, which can accommodate the Miniprep columns. The use of a vacuum regulator is also recommended. A negative pressure of -25-30 inches Hg is required. This is equivalent to approximately -850-1,000 mbar and -12-15 psi.

Lysis and homogenization using a mortar and pestle

1. Select 1-20 mg of animal tissue and transfer to a mortar, pre-chilled on ice. Grind rapidly and vigorously to form a homogenate.

Note: the following tissue-types should be completely frozen in liquid nitrogen before grinding:

- DNase-rich tissues, such as pancreas, thymus, lymphoid tissue, etc.
- Collagen-rich tissues, such as skin, connective tissue, etc.
- Keratoprotein-rich tissues or hard tissues, such as bone.

2. Add 350 µl of phosphate-buffered saline (PBS, not provided) and 0.9 µl of RNase A. Gently grind for 30 seconds to homogeneously mix the PBS with the ground tissue.

Note: For those tissue types in Step 1 (above) that require freezing in liquid nitrogen, please perform the following steps after pulverizing: Add 350 µl of PBS and 0.9 µl of RNase A. Warm the mortar to 56°C in the water bath until the PBS just melts. Gently grind for 1 minute. Then **proceed to Step 3, below.**

3. Collect 350 µl of the homogenate and transfer to a 2 ml microfuge tube (provided). If the volume of the homogenate is less than 350 µl, make it up to 350 µl with PBS.

4. Add 20 μ l Proteinase K and 150 μ l Buffer C-L. Mix immediately by vortexing for 1 minute.

Note: Do not add Proteinase K directly to buffer C-L.

5. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
6. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
7. Centrifuge for 10 minutes at 12,000xg at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

8. Attach the vacuum manifold base to a vacuum source. Firmly position the Miniprep column(s) into the complementary fittings on the manifold top. Transfer the clarified supernatant obtained from Step 7 to the Miniprep column. Turn on the vacuum source and adjust to -25 inches Hg. Continue to apply the vacuum until no solution remains in the Miniprep column.
9. Add 500 μ l of Buffer W1 and draw all of the solution through the column.
10. Add 700 μ l of Buffer W2 along the wall of Miniprep column to wash off residual Buffer W1 and draw all of the solution through the column. Repeat this wash step with a second 700 μ l aliquot of Buffer W2.

Note: Make sure that ethanol has been added into Buffer W2 concentrate.

Note: Add Buffer W2 along the tube wall to wash off any residual salt.

Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.

11. Transfer the Miniprep column to a 2 ml microfuge tube (provided) and centrifuge for 1 minute at 12,000xg.
12. Transfer the Miniprep column to a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 μ l of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.

Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

[Centrifuge protocol]

Lysis and homogenization using a mortar and pestle

1. Select 1-20 mg of animal tissue and transfer to a mortar, pre-chilled on ice. Grind rapidly and vigorously to form a homogenate.

Note: the following tissue-types should be completely frozen in liquid nitrogen before grinding:

- DNase-rich tissues, such as pancreas, thymus, lymphoid tissue, etc.
- Collagen-rich tissues, such as skin, connective tissue, etc.
- Keratin-rich tissues or hard tissues, such as bone.

2. Add 350 μ l of PBS (not provided), and 0.9 μ l of RNase A. Gently grind for 30 seconds to homogeneously mix the PBS with the ground tissue.

Note: For those tissue types in Step 1 (above) requiring freezing in liquid nitrogen, please perform the following steps after pulverizing: Add 350 μ l of PBS and 0.9 μ l of RNase A. Warm the mortar to 56°C in water bath until the PBS just melts. Gently grind for 1 minute. Then **proceed to Step 3, below**

3. Collect 350 μ l of the homogenate and transfer to a 2 ml microfuge tube (provided). If the volume of the homogenate is less than 350 μ l, make it up to 350 μ l with PBS.
4. Add 20 μ l Proteinase K and 150 μ l Buffer C-L. Mix immediately by vortexing for 1 minute.
Note: Do not add Proteinase K directly to buffer C-L.
5. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
6. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
7. Centrifuge at 12,000xg for 10 minutes at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

8. Place a Miniprep column into a 2 ml microfuge tube (provided). Pipette the clarified supernatant obtained from step 7 into the Miniprep column. Centrifuge for 1 minute at 12,000xg.
9. Discard the filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Pipette 500 μ l of Buffer W1 to the Miniprep column and centrifuge at 12,000xg for 1 minute.
10. Discard the filtrate and place the Miniprep column back into the 2 ml microfuge tube. Add 700 μ l of Buffer W2 and centrifuge for 1 minute at 12,000xg. Discard the filtrate from the 2 ml microfuge tube and repeat this wash step with a second 700 μ l aliquot of Buffer W2.
Note: Make sure that ethanol has been added into Buffer W2 concentrate.
Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.
11. Discard filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube and centrifuge for 1 minute at 12,000xg.
12. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 μ l of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.
Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

II. Purification of Genomic DNA from Plant Tissues and Filamentous Fungi

[Vacuum Protocol]

This procedure requires the use of AxyVac vacuum manifold or other manifold with complementary luer fittings which can accommodate the Miniprep columns. The use of a vacuum regulator is also recommended. A negative pressure of -25-30 inches Hg is required. This is equivalent to approximately -850-1,000 mbar and -12-15 psi.

Lysis and homogenization using a mortar and pestle

1. Using Table 1 (below), weigh out the appropriate amount of either fresh plant tissue or filamentous fungi and transfer to a mortar. The amount should be reduced by half if lyophilized, dehydrated, or dried tissues are used. Carefully add liquid nitrogen directly to the sample until it

is completely frozen. Use the pestle to pulverize it quickly and vigorously until it is reduced to a fine powder.

Table 1. Types of fresh plant tissues and filamentous fungi used for genomic DNA preparation

Flower or leaves	10-100 mg
Plant stem	≤ 240 mg
Plant root	≤ 240 mg
Plant seed	≤ 240 mg
Filamentous fungi	5-50 mg

Note: If cultured plant cells are used, collect 2×10^3 - 1×10^7 plant cells and spin for 1 minute at 10,000xg to pellet the cells. Resuspend the plant cells in 150 μ l of PBS and transfer to the mortar. Carefully add liquid nitrogen directly to the sample until it is completely frozen. Use the pestle to pulverize it quickly and vigorously until it is reduced to a fine powder. Add liquid nitrogen as required to prevent the material from thawing during pulverization. After pulverizing, warm the mortar to 56°C in a water bath until the pulverized material just melts. **Proceed to Step 2, below.**

2. Add 350 μ l of PBS (not provided), and 0.9 μ l of RNase A. Gently grind for 30 seconds to homogeneously mix the PBS with the ground tissue.

Note: For those tissue types in Step 1 (above) requiring freezing in liquid nitrogen, please perform the following steps after pulverizing: Add 350 μ l of PBS and 0.9 μ l of RNase A. Warm the mortar to 56°C in water bath until the PBS just melts. Gently grind for 1 minute. Then **proceed to Step 3, below.**

Note: Incomplete grinding will reduce the yield of genomic DNA.

Note: When the weight of the fresh plant tissue is >120 mg or the dried plant tissue is >60 mg, add 0.7 ml of PBS. After Step 2 has been completed, divide the sample evenly between two 2 ml microfuge tubes. Steps 4-7 will proceed in two parallel 2 ml microfuge tubes. In Step 8, the contents of the two tubes will be consolidated into a single Miniprep column.

3. Transfer the tissue homogenate into a 2 ml microfuge tube (provided). Determine the approximate volume. If the volume of the homogenate is less than 350 μ l, make it up to 350 μ l with PBS.
4. Add 20 μ l Proteinase K and 150 μ l Buffer C-L. Mix immediately by vortexing for 1 minute.

Note: Do not add Proteinase K directly to buffer C-L.

Note: If fibrous samples such as plant stem and root, or starch- and protein-rich samples such as seeds are used, increase the incubation time to 30 minutes in the water bath.

5. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from underside the lid.
6. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
7. Centrifuge at 12,000xg for 10 minutes at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

8. Attach the vacuum manifold base to a vacuum source. Firmly position the Miniprep column(s) into the complementary fittings on the manifold top. Transfer the clarified supernatant obtained from Step 7 to the Miniprep column. Turn on the vacuum source and adjust to -25 inches Hg. Continue to apply the vacuum until no solution remains in the Miniprep column.

9. Add 500 µl of Buffer W1. Draw all of the solution through the column.
10. Apply 700 µl of Buffer W2 along the wall of Miniprep column to wash off residual Buffer W1. Draw all of the solution through the column. Repeat this wash step with a second 700 µl aliquot of Buffer W2.

Note: Make sure that ethanol has been added into Buffer W2 concentrate.

Note: Add Buffer W2 along the tube wall to wash off any residual salt.

Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.

11. Transfer the Miniprep column into a 2 ml microfuge tube (provided) and centrifuge for 1 minute at 12,000xg.
12. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 µl of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.

Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

[Centrifuge Protocol]

Lysis and homogenization using a mortar and pestle

1. Using Table 1 (below), weigh out the appropriate amount of either fresh plant tissue or filamentous fungi and transfer to a mortar. The amount should be reduced by half if lyophilized, dehydrated, or dried tissues are used. Carefully add liquid nitrogen directly to the sample until it is completely frozen. Use the pestle to pulverize it quickly and vigorously until it is reduced to a fine powder.

Table 1. Types of fresh plant tissues and filamentous fungi used for genomic DNA preparation.

Flower or leaves	10-100 mg
Plant stem	≤ 240 mg
Plant root	≤ 240 mg
Plant seed	≤ 240 mg
Filamentous fungi	5-50 mg

Note: If cultured plant cells are used, collect 2×10^3 - 1×10^7 plant cells and spin for 1 minute at 10,000xg to pellet the cells. Resuspend the plant cells in 150 µl of PBS and transfer to the mortar. Carefully add liquid nitrogen directly to the sample until it is completely frozen. Use the pestle to pulverize it quickly and vigorously until it is reduced to a fine powder. Add liquid nitrogen as required to prevent the material from thawing when grinding. After pulverizing, warm the mortar to 56°C in a water bath until the pulverized material just melts. **Proceed to Step 2, below.**

2. Add 350 µl of PBS (not provided), and 0.9 µl of RNase A. Gently grind for 30 seconds to homogenously mix the PBS with the ground tissue.

Note: For those tissue types in Step 1 (above) requiring freezing in liquid nitrogen, please perform the following steps after pulverizing: Add 350 µl of PBS and 0.9 µl of RNase A. Warm the mortar to 56°C in water bath until the PBS just melts. Gently grind for 1 minute. Then **proceed to Step 3, below**

Note: Incomplete grinding will reduce the yield of genomic DNA.

Note: When the weight of the fresh plant tissue is >120 mg or the dried plant tissue is >60 mg, add 0.7 ml of PBS. After Step 2 has been completed, divide the sample evenly between two 2 ml microfuge tubes. Steps 4-7 will proceed in two parallel 2 ml microfuge tubes. In Step 8, the contents of the two tubes will be consolidated into a single Miniprep column.

3. Transfer the tissue homogenate into a 2 ml microfuge tube (provided). Determine the approximate volume. If the volume of the homogenate is less than 350 μ l, make it up to 350 μ l with PBS.
4. Add 20 μ l Proteinase K and 150 μ l Buffer C-L. Mix immediately by vortexing for 1 minute.
Note: Do not add Proteinase K directly to buffer C-L.
Note: If fibrous samples such as plant stem and root, or starch- and protein-rich samples such as seeds are used, increase the incubation time to 30 minutes in the water bath.
5. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
6. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
7. Centrifuge for 10 minutes at 12,000xg at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

8. Place a Miniprep column into a 2 ml microfuge tube (provided). Pipette the clarified supernatant obtained from step 7 into the Miniprep column. Centrifuge for 1 minute at 12,000xg.
9. Discard the filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Pipette 500 μ l of Buffer W1 to the Miniprep column and centrifuge for 1 minute at 12,000xg.
10. Discard the filtrate and place the Miniprep column back into the 2 ml microfuge tube. Add 700 μ l of Buffer W2 and centrifuge for 1 minute at 12,000xg. Discard the filtrate from the 2 ml microfuge tube and repeat this wash step with a second 700 μ l aliquot of Buffer W2.
Note: Make sure that ethanol has been added into Buffer W2 concentrate.
Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.
11. Discard filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Centrifuge for 1 minute at 12,000xg.
12. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 μ l of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.
Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

III. Purification of Genomic DNA from Cultured Animal Cells, Lymphocytes, Bone Marrow and Bone.

[Vacuum Protocol]

This procedure requires the use of AxyVac vacuum manifold or other manifold with complementary luer fittings which can accommodate the Miniprep columns. The use of a vacuum regulator is also recommended. A negative pressure of $-25-30$ inches Hg is required. This is equivalent to approximately $-850-1,000$ mbar and $-12-15$ psi.

Lysis and homogenization of sample

Select method A-E, depending upon the type of starting material used. If genomic DNA is extracted from plant cells, please follow the previous protocol "Purification of Genomic DNA from Plant Tissues" (above) to homogenize the plant cells.

A. Cells grown in suspension or a cell suspension freshly isolated from animal or human tissues:

- 1A. Collect 1×10^3 - 2×10^6 cells in suspension and transfer to a 2 ml microfuge tube (provided). Centrifuge for 5 minutes at 2,000xg to pellet the cells. Discard the supernatant.
- 2A. Add 350 μ l of PBS to resuspend the cells and then add 150 μ l of Buffer C-L. Let the tube stand for 1 minute at room temperature. **Proceed to Step 3, below.**

B. Cells grown in a monolayer in a 96-well, 24-well, 12-well or 6-well plate:

- 1B. Discard as much of the supernatant as possible, then add 350 μ l of PBS into each well and 150 μ l Buffer C-L. Let the plate stand for 1 minute at room temperature.
- 2B. Pipette up and down several times, and then transfer 500 μ l of the cell lysate to a 2 ml microfuge tube (provided).

C. Lymphocytes:

Suspend the lymphocytes in 250 μ l of PBS. **Proceed to Step 3, below.**

D. Bone marrow:

To obtain bone marrow samples (e.g., mouse), remove the femur and cut through the bone at both ends. Using a syringe and small gauge needle, inject 0.3 ml of PBS through one end of the bone and flush the marrow into a 2 ml microfuge tube (provided). Collect 250 μ l of marrow suspension. **Proceed to Step 3, below.**

E. Bone:

Either fresh or archaic bone can be processed using this method. The maximum amount of bone is 50 mg.

- 1E. Flash-freeze the bone sample by immersion in liquid nitrogen.
- 2E. Transfer the frozen bone to a mortar and pestle and grind to a fine powder. It may be necessary to add small amounts of liquid nitrogen to the mortar to keep the sample frozen for improved grinding.

Note: Maintaining the bone in a frozen state will improve grinding.

- 3E. Transfer the pulverized bone to a 2 ml microfuge tube (provided) and add 300 μ l of PBS. Vortex for 20 seconds.

- 4E. Place the microfuge tube on ice for 10 minutes. Allow the pulverized bone to settle to the bottom of the tube.
- 5E. Aspirate off 250 μ l of PBS. **Proceed to Step 3, below.**
3. Add 0.8 μ l of RNase A and 8 μ l Proteinase K. Vortex for 15 seconds and Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
- Note:** Do not add Proteinase K directly to buffer C-L.
4. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
5. Centrifuge for 10 minutes at 12,000xg at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

6. Attach the vacuum manifold base to a vacuum source. Firmly position the Miniprep column(s) into the complementary fittings on the manifold top. Transfer the clarified supernatant obtained from Step 5 to the Miniprep column. Turn on the vacuum source and adjust to -25 inches Hg. Continue to apply the vacuum until no solution remains in the Miniprep column.
7. Add 500 μ l of Buffer W1. Draw all of the solution through the column.
8. Add 700 μ l of Buffer W2 along the wall of Miniprep column to wash off residual Buffer W1, draw all of the solution through the column. Repeat this wash step with a second 700 μ l aliquot of Buffer W2.
- Note:** Make sure that ethanol has been added into Buffer W2 concentrate.
- Note:** Add Buffer W2 along the tube wall to wash off any residual salt.
- Note:** Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.
9. Transfer the Miniprep column into a 2 ml microfuge tube (provided) and centrifuge for 1 minute at 12,000xg.
10. Transfer the Miniprep column to a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 μ l of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.
- Note:** Pre-warming water or Eluent at 65°C will often improve elution efficiency.

[Centrifuge Protocol]

Lysis and homogenization of sample

Select method A-E, depending upon the type of starting material used. If genomic DNA is extracted from plant cells, please follow the previous protocol "Purification of Genomic DNA from Plant Tissues" (above) to homogenize the plant cells.

A. Cells grown in suspension or a cell suspension freshly-isolated from animal or human tissues:

- 1A. Collect 1×10^3 - 2×10^6 cells in suspension and transfer to a 2 ml microfuge tube (provided). Centrifuge for 5 minutes at 2,000xg to pellet the cells. Discard the supernatant.

2A. Add 350 μ l of PBS to resuspend the cells and then add 150 μ l of Buffer C-L. Let the tube stand for 1 minute at room temperature. **Proceed to Step 3, below.**

B. Cells grown in a monolayer in a 96-well, 24-well, 12-well or 6-well plate:

1B. Discard as much of the supernatant as possible, then add 350 μ l of PBS into each well and 150 μ l Buffer C-L. Let the plate stand for 1 minute at room temperature.

2B. Pipette up and down several times, then transfer 500 μ l of the cell lysate to a 2 ml microfuge tube (provided).

C. Lymphocytes:

Suspend the lymphocytes in 250 μ l of PBS. **Proceed to Step 3, below.**

D. Bone marrow:

To obtain bone marrow samples (e.g., mouse), remove the femur and cut through the bone at both ends. Using a syringe and small gauge needle, inject 0.3 ml of PBS through one end of the bone and flush the marrow into a 2 ml microfuge tube (provided). Collect 250 μ l of marrow suspension. **Proceed to Step 3, below.**

E. Bone:

Either fresh or archaic bone can be processed using this method. The maximum amount of bone is 50 mg.

1E. Flash-freeze the bone sample by immersion in liquid nitrogen.

2E. Transfer the frozen bone to a mortar and pestle and grind to a fine powder. It may be necessary to add small amounts of liquid nitrogen to the mortar to keep the sample frozen for improved grinding.

Note: Maintaining the bone in a frozen state will improve grinding.

3E. Transfer the pulverized bone to a 2 ml microfuge tube (provided) and add 300 μ l of PBS. Vortex for 20 seconds.

4E. Place the microfuge tube on ice for 10 minutes. Allow the pulverized bone to settle to the bottom of the tube.

5E. Aspirate off 250 μ l of PBS. **Proceed to Step 3, below.**

3. Add 0.8 μ l of RNase A and 8 μ l Proteinase K. Vortex for 15 seconds and incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.

Note: Do not add Proteinase K directly to buffer C-L.

4. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.

5. Centrifuge for 10 minutes at 12,000xg at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

6. Place a Miniprep column into a 2 ml microfuge tube (provided). Pipette the clarified supernatant obtained from step 5 into the Miniprep column. Centrifuge for 1 minute at 12,000xg.

7. Discard the filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Pipette 500 µl of Buffer W1 to the Miniprep column and centrifuge for 1 minute at 12,000xg.
8. Discard the filtrate and place the Miniprep column back into the 2 ml microfuge tube. Add 700 µl of Buffer W2 and centrifuge for 1 minute at 12,000xg. Discard the filtrate from the 2 ml microfuge tube and repeat this wash step with a second 700 µl aliquot of Buffer W2.
Note: Make sure that ethanol has been added into Buffer W2 concentrate.
Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.
9. Discard filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Centrifuge for 1 minute at 12,000xg.
10. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 µl of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.
Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

IV. Purification of Genomic DNA from Buccal Swabs

[Vacuum protocol]

This procedure requires the use of AxyVac vacuum manifold or other manifold with complementary luer fittings which can accommodate the Miniprep columns. The use of a vacuum regulator is also recommended. A negative pressure of -25-30 inches Hg is required. This is equivalent to approximately -850-1,000 mbar and -12-15 psi.

Lysis and homogenization of sample

1. Add 350 µl of PBS to a clean 2 ml microfuge tube (provided). Dip the buccal swab into the PBS and rotate against the wall of the tube for 30 seconds. Lift the swab above the PBS and rotate against the wall of the tube to wring out residual buffer and cells. Discard the swab. Add 1 µl of RNase A and mix briefly.
2. Add 8 µl Proteinase K, followed by 150 µl Buffer C-L to the sample. Mix immediately by vortex 30 Seconds
Note: Do not add Proteinase K directly to Buffer C-L.
3. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
4. Add 350 µl Buffer P-D to the sample, vortex for mix well and centrifuge for 10 minutes at 12,000xg.

Binding, washing and elution on the Miniprep column

5. Attach the vacuum manifold base to a vacuum source. Firmly position the Miniprep column(s) into the complimentary fittings on the manifold top. Transfer the clarified supernatant obtained

from Step 4 to the Miniprep column. Turn on the vacuum source and adjust to -25 inches Hg. Continue to apply the vacuum until no solution remains in the Miniprep column.

6. Add 500 μ l of Buffer W1 and draw all of the solution through the column.
7. Add 700 μ l of Buffer W2 along the wall of Miniprep column to wash off residual Buffer W1 and draw all of the solution through the column. Repeat this wash with a second 700 μ l aliquot of Buffer W2.

Note: Make sure that ethanol has been added into Buffer W2 concentrate.

Note: Add Buffer W2 along the tube wall to wash off any residual salt.

Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.

8. Transfer the Miniprep column into a 2 ml microfuge tube (provided) and centrifuge for 1 minute at 12,000xg.
9. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 μ l of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.

Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

[Centrifuge protocol]

Lysis and homogenization of sample

1. Add 350 μ l of PBS to a clean 2 ml microfuge tube (provided). Dip the buccal swab into the PBS and rotate against the wall of the tube for 30 seconds. Lift the swab above the PBS and rotate against the wall of the tube to wring out residual buffer and cells. Discard the swab. Add 1 μ l of RNase A and mix briefly.
2. Add 8 μ l of Proteinase K, followed by 150 μ l Buffer C-L to the sample. Mix immediately by vortexing for 30 seconds.

Note: Do not add Proteinase K directly to Buffer C-L.

3. Incubate at 56°C for 15 minutes. Following this, centrifuge briefly to remove condensation from inside the lid.
4. Add 350 μ l Buffer P-D to the sample. Vortex for 30 seconds and centrifuge for 10 minutes at 12,000xg.

Binding, washing and elution on the Miniprep column

5. Place a Miniprep column into a 2 ml microfuge tube (provided). Pipette the clarified supernatant obtained from step 4 into the Miniprep column. Centrifuge for 1 minute at 12,000xg.
6. Discard the filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Add 500 μ l of Buffer W1 to the Miniprep column and centrifuge for 1 minute at 12,000xg.

7. Discard the filtrate and place the Miniprep column back into the 2 ml microfuge tube. Add 700 μ l of Buffer W2 and centrifuge for 1 minute at 12,000xg. Discard the filtrate from the 2 ml microfuge tube and repeat this wash with a second 700 μ l aliquot of Buffer W2.

Note: Make sure that ethanol has been added into Buffer W2 concentrate.

Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.

8. Discard the filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Centrifuge for 1 minute at 12,000xg to remove residual W2.
9. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 μ l of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.

Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

V. Purification of Genomic DNA from Yeast

[Vacuum Protocol]

This procedure requires the use of AxyVac vacuum manifold or other manifold with complementary luer fittings which can accommodate the Miniprep columns. The use of a vacuum regulator is also recommended. A negative pressure of -25-30 inches Hg is required. This is equivalent to approximately -850-1,000 mbar and -12-15 psi.

Lysis and homogenization using a mortar and pestle

1. Collect 2×10^6 - 5×10^7 yeast cells and centrifuge for 1 minute at 10,000xg to pellet the cells. Resuspend the yeast cells in 350 μ l of PBS and transfer to a mortar.

Note: For yeast, an $OD_{600} = 1$ is approximately 3×10^7 cells/ml.

2. Gradually add liquid nitrogen until the yeast suspension is completely frozen. Using the pestle, quickly and forcefully reduce it to a fine powder. Add liquid nitrogen to prevent the sample from thawing during pulverization. After grinding is complete, warm the mortar at 56°C in water bath until it just begins to melt.
3. Add 150 μ l of Buffer C-L and 1.2 μ l of RNase A. Quickly grind the sample for 30 seconds.
4. Transfer 500 μ l of the yeast homogenate to a 2 ml microfuge tube (provided). If the volume of the homogenate is less than 500 μ l, add additional Buffer C-L up to 500 μ l.
5. Add 20 μ l Proteinase K. Mix immediately by vortexing for 30 seconds.

Note: Do not add Proteinase K directly to buffer C-L.

6. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
7. Add 350 μ l Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
8. Centrifuge for 10 minutes at 12,000xg at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

9. Attach the vacuum manifold base to a vacuum source. Firmly position the Miniprep column(s) into the complementary fittings on the manifold top. Transfer the clarified supernatant obtained from Step 8 to the Miniprep column. Turn on the vacuum source and adjust to -25 inches Hg. Continue to apply the vacuum until no solution remains in the Miniprep column.
10. Add 500 µl of Buffer W1. Draw all of the solution through the column.
11. Add 700 µl of Buffer W2 along the wall of Miniprep column to wash off residual Buffer W1, draw all of the solution through the column. Repeat this wash step with a second 700 µl aliquot of Buffer W2.
Note: Make sure that ethanol has been added into Buffer W2 concentrate.
Note: Add Buffer W2 along the tube wall to wash off any residual salt.
Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.
12. Transfer the Miniprep column into a 2 ml microfuge tube (provided) and centrifuge for 1 minute at 12,000xg.
13. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 µl of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.
Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

[Centrifuge Protocol]**Lysis and homogenization using a mortar and pestle**

1. Collect 2×10^6 - 5×10^7 yeast cells and centrifuge for 1 minute at 10,000xg to pellet the cells. Resuspend the yeast cells in 350 µl of PBS and transfer to a mortar.
Note: For yeast, an $OD_{600} = 1$ is approximately 3×10^7 cells/ml.
2. Gradually add liquid nitrogen until the yeast suspension is completely frozen. Using the pestle, quickly and forcefully reduce it to a fine powder. Add liquid nitrogen to prevent the sample from thawing during pulverization. After grinding is complete, warm the mortar at 56°C in water bath until it just begins to melt.
3. Add 150 µl of Buffer C-L and 1.2 µl of RNase A. Quickly grind the sample for 30 seconds.
4. Transfer 500 µl of the yeast homogenate into a 2 ml microfuge tube (provided). If the volume of the homogenate is less than 500 µl, add additional Buffer C-L up to 500 µl.
5. Add 20 µl Proteinase K. Mix immediately by vortexing for 30 seconds.
Note: Do not add Proteinase K directly to buffer C-L.
6. Incubate at 56°C for 15 minutes. Briefly centrifuge to remove drops from inside the lid.
7. Add 350 µl Buffer P-D to the sample and mix by vortexing at top speed for 30 seconds.
8. Centrifuge for 10 minutes at 12,000xg at ambient temperature to pellet cellular debris.

Binding, washing and elution on the Miniprep column

9. Place a Miniprep column to a 2 ml microfuge tube (provided). Pipette the clarified supernatant obtained from step 8 into the Miniprep column. Centrifuge for 1 minute at 12,000xg.
10. Discard the filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Add 500 µl of Buffer W1 to the Miniprep column and centrifuge for 1 minute at 12,000xg.
11. Discard the filtrate and place the Miniprep column back into the 2 ml microfuge tube. Add 700 µl of Buffer W2 and centrifuge for 1 minute at 12,000xg. Discard the filtrate from the 2 ml microfuge tube and repeat this wash step with a second 700 µl aliquot of Buffer W2.

Note: Make sure that ethanol has been added into Buffer W2 concentrate.

Note: Two washes with Buffer W2 are used to ensure the complete removal of salt, eliminating potential problems in subsequent enzymatic reactions.

12. Discard filtrate from the 2 ml microfuge tube. Place the Miniprep column back into the 2 ml microfuge tube. Centrifuge for 1 minute at 12,000xg.
13. Transfer the Miniprep column into a clean 1.5 ml microfuge tube (provided). To elute the genomic DNA, add 100-200 µl of Eluent (or deionized water) to the center of the membrane. Let it stand for 1 minute at room temperature. Centrifuge for 1 minute at 12,000xg.

Note: Pre-warming water or Eluent at 65°C will often improve elution efficiency.

Overview

Add 350 μ l of PBS
Add RNase A according to relevant protocols




Starting material

Add Proteinase K according to relevant protocols
Add 150 μ l of Buffer C-L
Incubate at 56°C for 15 minutes




Proteinase K digest

Add 350 μ l of Buffer P-D



Protein precipitation

Add 500 μ l of Buffer W1
Add 700 μ l of Buffer W2
Add 700 μ l of Buffer W2



Binding

Washing

Add 100-200 μ l of Eluent



Elution

Troubleshooting:

1. Low or no yield

- Insufficient starting material processed
- Inefficient lysis
- DNA not efficiently eluted
- Miniprep column membrane overdried during vacuum removal of Buffer W2

2. Low $A_{260/280}$

- Excessive material processed
- Inefficient lysis
- Contamination with protein

3. RNA present (elevated $A_{260/280}$)

- Failure to add RNase A.

4. Genomic DNA appears to be degraded

- Depending upon the completeness of degradation, the genomic DNA will either appear as a smear or as a smear trailing in front of a high molecular weight band on an agarose gel. Since no physical measure used during the purification process is sufficient to cause any visually discernable degradation, the most likely source is enzymatic. Many cells and tissues, exhibit high levels of endonuclease activity. These starting materials must be lysed rapidly and completely to prevent substantial enzymatic degradation of the genomic DNA. Flash-freezing in liquid nitrogen immediately after harvesting may be necessary.

5. Genomic DNA performs poorly in enzymatic reactions

- Low DNA concentration
- Salt contamination: insufficient Buffer W1 removal
- Ethanol contamination: insufficient centrifugation to remove residual Buffer W2

6. Clogged spin-filter

- Excessive material processed
- Inefficient lysis

7. Clogged Miniprep column

- Excessive material processed
- Inefficient lysis

Warranty/Disclaimer

AxyPrep Kits are designed for R&D and general laboratory use only. Axygen Biosciences makes no claims regarding the performance of these kits for clinical or diagnostic applications. Axygen Biosciences warrants that this kit will perform as indicated for the specified application for a period of up to 12 months from the date of receipt when stored in the specified manner and used according to the instructions provided. In using this product, the customer agrees that Axygen Biosciences shall not be held liable for any direct or indirect damages, including, but not limited to, personal injury, property damage or lost profits (or other economic loss) resulting from the use or inability to use this product. In the event that this product fails to perform in the specified manner, remedial measures on the part of Axygen Biosciences shall be limited to the replacement of this product and will be implemented at the discretion of Axygen Biosciences.