



# CD127 Monoclonal Antibody (eBioRDR5), APC, eBioscience™

<b>Product Details</b>		
Size	100 Tests	
Species Reactivity	Human	
Published Species	Human	
Host/Isotype	Mouse / IgG1, kappa	
Recommended Isotype Control	Mouse IgG1 kappa Isotype Control (P3.6.2.8.1), APC, eBioscience™	
Class	Monoclonal	
Туре	Antibody	
Clone	eBioRDR5	
Conjugate	APC	
Form	Liquid	
Concentration	5 μL/Test	
Purification	Affinity chromatography	
Storage buffer	PBS, pH 7.2, with 0.1% gelatin, 0.2% BSA	
Contains	0.09% sodium azide	
Storage conditions	4° C, store in dark, DO NOT FREEZE!	
RRID	AB_1659670	

Applications	Tested Dilution	Publications
Immunohistochemistry (Frozen) (IHC (F))	-	1 Publication
Flow Cytometry (Flow)	5 μL (0.06 μg)/test	7 Publications

## **Product Specific Information**

Description: The eBioRDR5 monoclonal antibody reacts with human CD127 (Interleukin-7 Receptor alpha). CD127 complexes with CD132, also known as the common gamma chain (gamma c), to form the multi-functional IL-7 receptor (IL-7R). CD127 is a type I alycoprotein with a molecular weight of 75-80 kDa and is expressed by immature B cells through the early pre-B stage, by thymocytes during several stages of their development, and on most mature T cells, with transient down-regulation upon activation. Binding of IL-7 results in signal transduction which occurs through several tyrosine kinase pathways including the Jak /STAT pathway. IL-7 is indispensible for the development of lymphocytes, and the control of homeostatic proliferation of T-cells in the periphery. In addition, IL-7R signaling is know to be involved in the regulation of T cell receptor (TCR) locus rearrangement in gamma delta T cells.

Interestingly, recently it has been demonstrated that CD127 expression is down-regulated on CD4+CD25+ regulatory T cells (T regs). While the co-expression of CD4 and CD25 has become widely used as an indicator of T regs, this method of identification may also include cells without suppressive activity. It has clearly been shown that CD4+CD25+ cells that have down-regulated the expression of CD127 are significantly more highly-enriched for the regulatory T population, as defined by expression of the T regspecific transcription factor Foxp3 and the suppressive activity of these cells, in vitro.

Binding of the eBioRDR5 monoclonal antibody to PBMCs is blocked by pre-incubation of the cells with recombinant human IL-7 (cat. 14-8079).


Applications Reported: This eBioRDR5 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This eBioRDR5 antibody has been pre-titrated and tested by flow cytometric analysis of normal human peripheral blood cells. This can be used at 5  $\mu$ L (0.06  $\mu$ g) per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test.

Excitation: 633-647 nm; Emission: 660 nm; Laser: Red Laser.

Filtration: 0.2 µm post-manufacturing filtered.

# Product Images For CD127 Monoclonal Antibody (eBioRDR5), APC, eBioscience™



## CD127 Antibody (17-1278-42) in Flow

Staining of normal human peripheral blood cells with Anti-Human CD3 FITC (Product # 11-0038-42) and Mouse IgG1 K Isotype Control APC (Product # 17-4714-81) (left) or Anti-Human CD127 APC (right). Cells in the lymphocyte gate were used for analysis.

### ■ 8 References

# Immunohistochemistry (Frozen) (1)

Journal of immunology (Baltimore, Md.: 1950)

Loss of IL-7 receptor alpha on CD4+ T cells defines terminally differentiated B cell-helping effector T cells in a B cell-rich lymphoid tissue.

Authors: Lim HW,Kim CH

Species Not Applicable

**Dilution** Not Cited

**Year** 2007

# Flow Cytometry (7)

#### Frontiers in immunology

Th2 Biased Immunity With Altered B Cell Profiles in Circulation of Patients With Sporotrichosis Caused by *Sporothrix globosa*.

"17-1278 was used in Flow cytometry/Cell sorting to shed light on the potential involvement of peripheral T and B cell immunity against this mycotic infection."

Authors: Zu J, Yao L, Song Y, Cui Y, Guan M, Chen R, Zhen Y, Li S

Species Human

**Dilution** Not Cited

**Year** 2021

#### Cancers

Murlentamab, a Low Fucosylated Anti-Müllerian Hormone Type II Receptor (AMHRII) Antibody, Exhibits Anti-Tumor Activity through Tumor-Associated Macrophage Reprogrammation and T Cell Activation.

"17-1278 was used in Flow cytometry/Cell sorting to demonstrate here that the murlentamab opsonization of AMHRII-expressing ovarian tumor cells, in the presence of unstimulated- or tumor-associated macrophage (TAM)-like macrophages, significantly promotes macrophage-mediated ADCC and shifts the whole microenvironment towards a pro-inflammatory and anti-tumoral status, thus triggering anti-tumor activity."

Authors: Prat M,Salon M,Allain T,Dubreuil O,Noël G,Preisser L,Jean B,Cassard L,Lemée F,Tabah-Fish I,Pipy B, Jeannin P,Prost JF,Barret JM,Coste A

Species Human

**Dilution** Not Cited

**Year** 2021

View more Flow references on thermofisher.com

# More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production of specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, his warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended uses not stended to anyone other than the Buyer. Any model or sample furnished to guerral substantially in the provided in the provided provided is subjected to provide and does not represent that any Product will conform to such model or sample. No OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES of MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPEAR, REPLACEMENT OR REPLAND FOR THE NON-CONFORMING PRODUCTS (S) AT SELLER'S SOLE OPTION. THERE IS NO DOBIGATION TO NEED, OR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses, or any type of consumption to human or animals.