

TLR4/MD-2 Complex Monoclonal Antibody (MTS510), PE, eBioscience™

| Product Details | |
|-----------------------------|---|
| Size | 100 µg |
| Species Reactivity | Mouse |
| Published Species | Rat, Mouse, Human |
| Host/Isotype | Rat / IgG2a, kappa |
| Recommended Isotype Control | Rat IgG2a kappa Isotype Control (eBR2a), PE, eBioscience™ |
| Class | Monoclonal |
| Type | Antibody |
| Clone | MTS510 |
| Conjugate | PE |
| Form | Liquid |
| Concentration | 0.2 mg/mL |
| Purification | Affinity chromatography |
| Storage buffer | PBS, pH 7.2, with 0.1% gelatin |
| Contains | 0.09% sodium azide |
| Storage conditions | 4° C, store in dark, DO NOT FREEZE! |
| RRID | AB_466264 |

| Applications | Tested Dilution | Publications |
|---|-----------------|-----------------|
| Immunohistochemistry (Frozen) (IHC (F)) | - | 1 Publication |
| Immunocytochemistry (ICC/IF) | - | 1 Publication |
| Flow Cytometry (Flow) | 0.5 µg/test | 27 Publications |

Product Specific Information

Description: The MTS510 monoclonal antibody reacts with the mouse Toll-like receptor 4 (TLR4)/MD-2 complex. At least ten members of the Toll family have been identified. This family of type I transmembrane proteins is characterized by an extracellular domain with leucine-rich repeats and a cytoplasmic domain with homology to the type I IL-1 receptor. Two of these receptors, TLR2 and TLR4, are pattern recognition receptors and signaling molecules in response to bacterial lipoproteins and have been implicated in innate immunity and inflammation. TLR4 physically associates with MD-2, and together with CD14, this complex is responsible for LPS recognition and signaling. In the mouse, TLR4 is expressed by thioglycolate-elicited peritoneal macrophages. Incubation of peritoneal macrophages with LPS results in down regulation of surface TLR4/MD-2. The TLR4 gene is defective in C3H/HeJ and C57BL/10ScCr mice, both of which have been well characterized as hyporesponders to LPS.

The MTS510 monoclonal antibody co-immunoprecipitates MD-2 (~30 kDa) and TLR4 (~100 kDa), and preferentially reacts with TLR4 that is associated with MD-2. In comparison, binding of the UT41 monoclonal antibody occurs with and without formation of the TLR4/MD-2 complex. Please contact eBioscience Technical Support for further information.


Applications Reported: MTS510 has been reported for use in flow cytometric analysis.

Applications Tested: The MTS510 antibody has been tested by flow cytometric analysis of mouse thioglycolate-elicited peritoneal exudate cells. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells /test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Excitation: 488-561 nm; Emission: 578 nm; Laser: Blue Laser, Green Laser, Yellow-Green Laser.

Filtration: 0.2 µm post-manufacturing filtered.

Product Images For TLR4/MD-2 Complex Monoclonal Antibody (MTS510), PE, eBioscience™



TLR4/MD-2 Complex Antibody (12-9924-82) in Flow

Staining of 3-day thioglycolate-induced BALB/c macrophages with 0.25 µg of Rat IgG2a kappa Isotype Control PE (Product # 12-4321-80) (open histogram) or 0.25 µg of Anti-Mouse TLR4/MD-2 Complex PE (filled histogram). Cells were pre-incubated with Anti-Mouse CD16/CD32 Purified (Product # 14-0161-82). Total viable cells were used for analysis.

[View more figures on thermofisher.com](http://thermofisher.com)

Immunohistochemistry (Frozen) (1)

Journal of immunology (Baltimore, Md. : 1950)

TLRs regulate the gatekeeping functions of the intestinal follicle-associated epithelium.

Authors: Chabot S,Wagner JS,Farrant S,Neutra MR

Species
Not Applicable

Dilution
Not Cited

Year
2006

Immunocytochemistry (1)

Scientific reports

IL-33/ST2 signalling and crosstalk with FcRI and TLR4 is targeted by the parasitic worm product, ES-62.

"12-9924 was used in Flow cytometry/Cell sorting to investigate whether ES62 targets the MyD88-IL-33 signalling network by subverting ST2-signalling to exhibit therapeutic potential in allergy."

Authors: Ball DH,AI-Riyami L,Harnett W,Harnett MM

Species
Mouse

Dilution
Not Cited

Year
2018

Flow Cytometry (27)

Journal of the American Heart Association

Sphingomyelin Synthase 2 Inhibition Ameliorates Cerebral Ischemic Reperfusion Injury Through Reducing the Recruitment of Toll-Like Receptor 4 to Lipid Rafts.

"12-9924 was used in Flow cytometry/Cell sorting to investigate the role of SMS2 in a mouse model of cerebral I/R."

Authors: Xue J,Yu Y,Zhang X,Zhang C,Zhao Y,Liu B,Zhang L,Wang L,Chen R,Gao X,Jiao P,Song G,Jiang XC,Qin S

Species
Mouse

Dilution
1:40

Year
2019

Immunity

Chanzyne TRPM7 Mediates the Ca²⁺ Influx Essential for Lipopolysaccharide-Induced Toll-Like Receptor 4 Endocytosis and Macrophage Activation.

"Published figure using TLR4/MD-2 Complex monoclonal antibody (Product # 12-9924-82) in Flow Cytometry"

Authors: Schappe MS,Szteyn K,Stremaska ME,Mendu SK,Downs TK,Seegren PV,Mahoney MA,Dixit S,Krupa JK,Stipes EJ,Rogers JS,Adamson SE,Leitinger N,Desai BN

Species
Not Applicable

Dilution
Not Cited

Year
2018

[View more Flow references on thermofisher.com](#)

More applications with references on thermofisher.com

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.