Performance guarenteed

# GL7 Monoclonal Antibody (GL-7 (GL7)), Alexa Fluor™ 488, eBioscience™

Product Details	
Size	100 µg
Species Reactivity	Human, Mouse
Published Species	Mouse, Human
Host/Isotype	Rat / IgM
Recommended Isotype Control	Rat IgM Isotype Control (eBRM), Alexa Fluor™ 488, eBioscience™
Class	Monoclonal
Туре	Antibody
Clone	GL-7 (GL7)
Conjugate	Alexa Fluor™ 488
Form	Liquid
Concentration	0.5 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2, with 0.1% gelatin
Contains	0.09% sodium azide
Storage conditions	4° C, store in dark, DO NOT FREEZE!
RRID	AB_2016717

Applications	Tested Dilution	Publications
Immunohistochemistry (IHC)	-	4 Publications
Immunocytochemistry (ICC/IF)	-	2 Publications
Flow Cytometry (Flow)	0.5 µg/test	18 Publications
Functional Assay (FN)	-	1 Publication

#### **Product Specific Information**

Description: This GL7 monoclonal antibody reacts with a cell-surface protein found on T and B lymphocytes activated in vitro, on bone marrow pre-B-II cells, germinal center B cells, and also human B cell lines Ramos and Daudi. There is strain variability with respect to antigen distribution on thymocytes and Con A-activated spleen cells, with expression in BALB/c greater than that in C57BL/6. GL7 is commonly used as a marker for mouse germinal center B cells. The epitope of GL7 has been identified as a sialic acid glycan moiety called Neu5Ac. This moiety is recognized by CD22.

Applications Reported: This GL-7 (GL7) antibody has been reported for use in flow cytometric analysis.


Applications Tested: This GL-7 (GL7) antibody has been tested by flow cytometric analysis of stimulated mouse splenocytes. This can be used at less than or equal to 0.5  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

1

Excitation: 488 nm; Emission: 519 nm; Laser: Blue Laser.

Filtration: 0.2 µm post-manufacturing filtered.

## Product Images For GL7 Monoclonal Antibody (GL-7 (GL7)), Alexa Fluor™ 488, eBioscience™



#### GL7 Antibody (53-5902-82) in Flow

Staining of unstimulated (left) or Con A-stimulated (right) C57Bl/6 splenocytes with Anti-Mouse CD3e PE (Product # 12-0031-82) and 0.25 µg of Anti-Human/Mouse GL7 (T and B Cell Activation Marker) Alexa Fluor® 488. Total viable cells, as determined by Fixable Viability Dye eFluor® 780 (Product # 65-0865-14) were used for analysis.

View more figures on thermofisher.com

### 25 References

### Immunohistochemistry (4)

Cell metabolism	<b>Species</b> Not Applicable
Microbiota-Derived Metabolites Suppress Arthritis by Amplifying Aryl-	
Hydrocarbon Receptor Activation in Regulatory B Cells.	Dilution
"Published figure using GL7 monoclonal antibody (Product # 53-5902-82) in Immunohistochemistry"	Not Cited
Authors: Rosser EC,Piper CJM,Matei DE,Blair PA,Rendeiro AF,Orford M,Alber DG,Krausgruber T,Catalan D,Klein N, Manson JJ,Drozdov I,Bock C,Wedderburn LR,Eaton S,Mauri C	<b>Year</b> 2020
Frontiers in immunology	Species
Deficiency in Calcium-Binding Protein S100A4 Impairs the Adjuvant	Not Applicable
Action of Cholera Toxin.	Dilution
"Published figure using GL7 monoclonal antibody (Product # 53-5902-82) in Immunofluorescence"	Not Cited
Authors: Sun JB,Holmgren J,Larena M,Terrinoni M,Fang Y,Bresnick AR,Xiang Z	Year
	2019

View more IHC references on thermofisher.com

#### Immunocytochemistry (2)

Nature communications	Species
B cell activation and plasma cell differentiation are inhibited by de novo	Not Applicable <b>Dilution</b> Not Cited
DNA methylation.	
"Published figure using GL7 monoclonal antibody (Product # 53-5902-82) in Immunofluorescence"	
Authors: Barwick BG,Scharer CD,Martinez RJ,Price MJ,Wein AN,Haines RR,Bally APR,Kohlmeier JE,Boss JM	<b>Year</b> 2018
Nature communications	Species
Limited clonal relatedness between gut IgA plasma cells and memory B	Not Applicable
cells after oral immunization.	Dilution
"Published figure using GL7 monoclonal antibody (Product # 53-5902-82) in Immunofluorescence"	Not Cited
Authors: Bemark M,Hazanov H,Strömberg A,Komban R,Holmqvist J,Köster S,Mattsson J,Sikora P,Mehr R,Lycke NY	Year
	2016

#### More applications with references on thermofisher.com

Flow (18) FN (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of sublishity for use in applications regulated by FDA is made. The warranty for value of shipment when the Product is subjected to normal, proper and introded usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent on anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTABILTY, FITNESS FOR ANY PARTICULOUTS DURING THE WARRANTER TO REPLANE FOR THE WONCONST DURING FOR DOBLIGATION TO REPAIR, REPLACE OR REPLANE FOR DESCHAPTING DISCHAPTING DISCHAPTING