Performance guarenteed

CD4 Monoclonal Antibody (GK1.5), NovaFluor Blue 555, eBioscience™

| Product Details | |
|--------------------|-------------------------------------|
| Size | 25 µg |
| Species Reactivity | Mouse |
| Host/Isotype | Rat / IgG2b, kappa |
| Class | Monoclonal |
| Туре | Antibody |
| Clone | GK1.5 |
| Conjugate | NovaFluor™ Blue 555 |
| Form | Liquid |
| Concentration | 0.1 mg/mL |
| Purification | Affinity chromatography |
| Storage buffer | PBS, pH 7.2 |
| Contains | 0.09% sodium azide |
| Storage conditions | 4° C, store in dark, DO NOT FREEZE! |

| Applications | Tested Dilution | Publications |
|-----------------------|-----------------|--------------|
| Flow Cytometry (Flow) | 0.2 µg/test | - |

Product Specific Information

Description: The GK1.5 monoclonal antibody reacts with the mouse CD4 molecule, a 55 kDa cell surface receptor expressed by a majority of thymocytes, subpopulation of mature T cells and dendritic cells. CD4 binds to MHC class II on the surface of antigen presenting cells and plays an important role both in T cell development and in optimal functioning of mature T cells. In T cells, CD4 associates with protein tyrosine kinase p56lck through its cytoplasmic tail. Binding of GK1.5 is blocked by RM4-5.

Applications Reported: This GK1.5 antibody has been reported for use in flow cytometric analysis.

Applications Tested: This GK1.5 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.2 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.


NovaFluor dyes are not compatible with DNA intercalating viability dyes. Do not use viability dyes such as propidium iodide, 7actinomycin D (7-AAD) and DAPI. Invitrogen LIVE/DEAD Fixable Dead Cell stains are recommended for use with NovaFluor dyes.

Each NovaFluor conjugate or kit is shipped with CellBlox Blocking Buffer. Use this buffer whenever staining with NovaFluor conjugates, including single-color compensation controls using cells. Use 5 µL of CellBlox Blocking Buffer per stained cell sample containing 10^3 to 10^8 cells.

Excitation: 496 nm; Emission: 555 nm; Laser: 488 nm (Blue) Laser

NovaFluor conjugates are based on Phiton[™] technology utilizing novel nucleic acid dye structures that allow for engineered fluorescent signatures with consideration for spillover and spread impacts. Learn more

Product Images For CD4 Monoclonal Antibody (GK1.5), NovaFluor Blue 555, eBioscience™



CD4 Antibody (M001T02B03) in Flow

C57BL/6 mouse splenocytes were either left unstained (blue histogram) or stained with 0.2 µg of CD4 Monoclonal Antibody, NovaFluor Blue 555 (Product # M001T02B03) (purple histogram) and acquired in the B3 channel on a 5-laser Cytek Aurora. Cells in the lymphocyte gate were used in the analysis.

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of a set, as set forth in the Production documentation, specifications and for accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTES OF MERCHANTABILITY, FITNESS FOR ANY PARTICUAR PURPOSE, OR REVIN FOR DOX THE WARRANTES TO REVINE FOR CONCORDING PRODUCTS SUCH STER WARRANTES OR REPLAY DERIOD IS LIMITED TO REPARE, REPLACE MENT OF OR REPLAY DOR OR REPLAY DERION THE WORKEN THE UN-CONFORMING PRODUCTS) AS THE CONCOLORITIS DURING THE WARRANTES TO REPLAY DERION DELEGATION TO REPARE, REPLACE OR REPLAY DERION REPLAY. SOLE OFTICS ON LONG THE WARRANTES OR REPLAY DERION TO REPARE, REPLACE OR REPLAY DERION THE NON-CONFORMING PRODUCTS) AS THE RESULTO OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MALUREE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (V) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product is indeed for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo o