


| Table of Contents | Page |
|--|------|
| 1. Description | 1 |
| 2. Overview of GeneBLAzer [®] GPCR Cell-based Assays | 2 |
| 3. Materials Supplied | 3 |
| 4. Materials Required | 4 |
| 4.1 Optional Equipment and Materials | 4 |
| 5. Detailed Cell Handling Procedures | 5 |
| 5.1 Thawing Method | 5 |
| 5.2 Propagation Method | 5 |
| 5.3 Freezing Method | 5 |
| 6. Media Requirements | 6 |
| 7. Transient Transfection Procedure and Agonist Test Assay | 7 |
| 7.2 Detailed Assay Protocol | 8 |
| 7.3 Creating a Stable Cell Line | 9 |
| 8. Parental Cell Negative Control Assay Procedure | 10 |
| 8.1 Quick Assay Reference Guides | 10 |
| 8.2 Detailed Assay Protocol | 10 |
| 9. Detection | 11 |
| 10. Data Analysis | 12 |
| 10.1 Background Subtraction and Ratio Calculation | 12 |
| 10.2 Visual Observation of Intracellular Beta-lactamase Activity Using LiveBLAzer [™] -FRET B/G Substrate (CCF4-AM) | 12 |
| 10. References | 13 |
| 11. Purchaser Notification | 14 |

1. Description

GeneBLAzer[®] G α 15-NFAT-*bla* CHO-K1 cells contain the promiscuous G protein G α 15 stably integrated into the GeneBLAzer[®] G α 15-NFAT-*bla* CHO-K1 cell line. CellSensor[®] NFAT-*bla* CHO-K1 contains a beta-lactamase reporter gene under control of a NFAT response element stably integrated into CHO-K1 cells.

The GeneBLAzer[®] G α 15-NFAT-*bla* CHO-K1 cell line is designed for the construction of Gi/o coupled GPCR cell-based assays. Using Lipofectamine[™] 2000, you transfect a plasmid containing your Gi/o GPCR of interest into this cell line, and the promiscuous G α 15 protein couples the Gi/o GPCR to the NFAT-*bla* reporter construct. The cell line has also been functionally validated for Z'-Factor and EC₅₀ concentrations of Thapsigargin.



| | |
|--------------------------------|---------|
| EC ₅₀ | 2.74 nM |
| Z'-factor at EC ₁₀₀ | 0.77 |

Dose response of G α 15-NFAT-*bla* CHO-K1 cells to Thapsigargin.


For Technical Support for this or other Drug Discovery Products, dial 760-603-7200, option 3, extension 40266

2. Overview of GeneBLAzer® GPCR Cell-based Assays

The GeneBLAzer® GPCR Cell-based Assay provides a highly accurate, sensitive, and easy-to-use method of monitoring cellular response to drug candidates or other stimuli (1). The core of the GeneBLAzer® assay technology is a beta-lactamase (*bla*) fluorescence resonance energy transfer (FRET) substrate that generates a ratiometric reporter response with minimal experimental noise. In addition to the dual-color (blue/green) readout of stimulated and unstimulated cells, this ratiometric method reduces the absolute and relative errors that can mask the underlying biological response of interest. Such errors include variations in cell number, transfection efficiency, substrate concentration, excitation path length, fluorescence detectors, and volume changes. The GeneBLAzer® GPCR assay technology has been proven in high-throughput screening (HTS) campaigns for a range of target classes, including G-protein coupled receptors (GPCRs) (2, 3), nuclear receptors (4-6) and kinase signaling pathways (7).

The GeneBLAzer® assay technology uses a mammalian-optimized *bla* reporter gene combined with a FRET-enabled substrate to provide reliable and sensitive detection in cells (1) (Figure 1). Cells are loaded with an engineered fluorescent substrate containing two fluorophores: coumarin and fluorescein. In the absence of *bla* expression, the substrate molecule remains intact. In this state, excitation of the coumarin results in fluorescence resonance energy transfer to the fluorescein moiety and emission of green fluorescent light. However, in the presence of *bla* expression, the substrate is cleaved separating the fluorophores and disrupting energy transfer. Excitation of the coumarin in the presence of *bla* enzyme activity results in a blue fluorescence signal.

Figure 1.



Fluorescent detection of beta-lactamase reporter gene response using a FRET-enabled substrate. After substrate loading, in the absence of beta-lactamase expression, cells appear green fluorescent. In the presence of beta-lactamase expression, the substrate is cleaved and cells appear blue fluorescent.

3. Materials Supplied

| Product: | Name | Size | Catalog # |
|---------------------------------------|--|--------|-----------|
| | GeneBLAzer® Gα15-NFAT-<i>bla</i> CHO-K1 cells Includes: <ul style="list-style-type: none">• GeneBLAzer® Gα15-NFAT-<i>bla</i> CHO-K1 cells (K1213)• Protocol• Certificate of Analysis | 1 tube | K1213 |
| Shipping Condition: | Dry ice | | |
| Storage Condition of Cells: | Liquid nitrogen. Immediately upon receipt, cells must be stored in liquid nitrogen or thawed for immediate use. Cells stored at -80°C can quickly lose viability. | | |
| Growth Properties of Cells: | Adherent | | |
| Cell Phenotype: | Epithelial | | |
| Selection Marker(s) for Cells: | Zeocin™ 100 µg/ml; Blasticidin 5 µg/ml; | | |
| Mycoplasma Testing: | Negative | | |
| BioSafety Level: | 2 | | |

For Technical Support for this or other Drug Discovery Products, dial 760-603-7200, option 3, extension 40266

4. Materials Required

| Media/Reagents | Recommended Source | Part # |
|--|--------------------|---|
| LiveBLAzer [™] -FRET B/G Loading Kit: LiveBLAzer [™] -FRET B/G Substrate (CCF4-AM) DMSO for Solution A Solution B Solution C | Invitrogen | K1427 (70 μ g) K1095 (200 μ g) K1096 (1 mg) K1030 (5 mg) |
| Solution D | Invitrogen | K1156 (1 ml) K1157 (25 ml) |
| Recovery [™] Cell Culture Freezing Medium | Invitrogen | 12648-010 |
| DMEM (high-glucose), with GlutaMAX [™] | Invitrogen | 10569-010 |
| DMSO | Fluka | 41647 |
| Fetal bovine serum (FBS), dialyzed, (DO NOT SUBSTITUTE!) | Invitrogen | 26400-036 |
| Non-essential amino acids (NEAA) | Invitrogen | 11140-050 |
| Penicillin/Streptomycin (antibiotics) | Invitrogen | 15140-122 |
| Phosphate-buffered saline without calcium and magnesium [PBS(-)] | Invitrogen | 14190-136 |
| HEPES (1 M, pH 7.3) | Invitrogen | 15630-080 |
| 0.05% Trypsin/EDTA | Invitrogen | 25300-054 |
| Thapsigargin | Calbiochem | 586005 |
| Zeocin [™] | Invitrogen | R250-01 |
| Blasticidin | Invitrogen | R210-01 |
| Lipofectamine [™] 2000 | Invitrogen | 11668-019 |
| pcDNA [™] 6.2-hygro-DEST Vector (or other vector that does not contain a Zeocin [™] or Blasticidin resistance gene) | Invitrogen | K1233 |
| Opti-MEM [®] I Medium | Invitrogen | 11058-021 |

| Consumables | Recommended Source | Part # |
|--|---------------------|--------|
| Black-wall, clear-bottom, 96-well assay plates (with low fluorescence background) | Corning | 3603 |
| Black-wall, clear-bottom, 384-well assay plates (with low fluorescence background) | Corning | 3712 |
| Compressed air | Various | --- |
| Equipment | Recommended Source | |
| Fluorescence plate reader with bottom-read capabilities | Various | |
| Filters if required for plate reader (see Section 7.4.1) | Chroma Technologies | |

4.1 Optional Equipment and Materials

- Epifluorescence- or fluorescence-equipped microscope, with appropriate filters
- Microplate centrifuge

For Technical Support for this or other Drug Discovery Products, dial 760-603-7200, option 3, extension 40266

5. Detailed Cell Handling Procedures

Note: Refer to **Section 6, Media Requirements** for specific media recipes.

5.1 Thawing Method

Note: Cells are shipped on dry ice and as such may require a short period of time prior to full recovery and normal growth.

1. Place 14 ml of Thawing Medium into a T75 flask.
2. Place the flask in a humidified 37°C/5% CO₂ incubator for 15 minutes to allow medium to equilibrate to the proper pH and temperature.
3. Remove the vial of cells to be thawed from liquid nitrogen and rapidly thaw by placing at 37°C in a water bath with gentle agitation for 1–2 minutes. Do not submerge vial in water.
4. Decontaminate the vial by wiping with 70% ethanol before opening in a Class II biological safety cabinet.
5. Transfer the vial contents drop-wise into 10 ml of Thawing Medium in a sterile 15-ml conical tube.
6. Centrifuge cells at 200 × g for 5 minutes.
7. Aspirate supernatant and resuspend the cell pellet in 1 ml of fresh Thawing Medium.
8. Transfer contents to the T75 tissue culture flask containing pre-equilibrated Thawing Medium and place flask in the humidified 37°C/5% CO₂ incubator.
9. At first passage, switch to Growth Medium.

5.2 Propagation Method

1. Passage or feed cells at least twice a week. Maintain cells between 5% and 95% confluence. Do not allow cells to reach confluence.
2. To passage cells, aspirate medium, rinse once in PBS, add 0.05% Trypsin/EDTA (3 ml for a T75 flask, 5 ml for a T175 flask, and 7 ml for T225 flask) and swirl to coat the cells evenly. Cells usually detach after ~2–5 minutes exposure to 0.05% Trypsin/EDTA. Add an equal volume of Growth Medium to inactivate the 0.05% Trypsin/EDTA.
3. Verify under a microscope that cells have detached and clumps have completely dispersed.
4. Centrifuge cells at 200 × g for 5 minutes and resuspend in Growth Medium.

5.3 Freezing Method

1. Harvest the cells as described in **Section 5.2** (above), Step 2. After detachment, count the cells, centrifuge cells at 200 × g for 5 minutes, and resuspend in 4°C Freeze Medium to a density of 2E6 cells/ml.
2. Dispense 1.0-ml aliquots into cryogenic vials.
3. Place in an insulated container for slow cooling and store overnight at -80°C.
4. Transfer to liquid nitrogen the next day for storage.

6. Media Requirements

Note: Unless otherwise stated, have all media and solutions at least at room temperature (we recommend 37°C for optimal performance) before adding to cells.

Note: Make **NO MEDIA SUBSTITUTIONS**, as these cell lines have been specifically validated for optimal assay performance with these media. For dividing cells, we recommend that you create and store an aliquot for back up.

| Component | Assay Medium | Growth Medium | Thawing Medium | Freeze Medium |
|--|----------------|----------------|----------------|---------------|
| DMEM | 90% | 90% | 90% | — |
| Dialyzed FBS (Do not substitute!) | 10% | 10% | 10% | — |
| NEAA | 0.1 mM | 0.1 mM | 0.1 mM | — |
| HEPES (pH 7.3) | 25 mM | 25 mM | 25 mM | — |
| Penicillin (antibiotic) | 100 U/ml | 100 U/ml | 100 U/ml | — |
| Streptomycin (antibiotic) | 100 μ g/ml | 100 μ g/ml | 100 μ g/ml | — |
| Recovery™ Cell Culture Freezing Medium | — | — | — | 100% |
| Zeocin™ | — | 100 μ g/ml | — | — |
| Blasticidin | — | 5 μ g/ml | — | — |

7. Transient Transfection Procedure and Agonist Test Assay

The following procedure describes how to transfect your Gi/o GPCR of interest into this G α 15-containing cell line and then perform an assay using a known agonist to test for a positive beta-lactamase response from the NFAT response element. The parental cell line should be tested separately for any endogenous response to the agonist, as described in **Section 8**.

Note: G α 15 may not couple to all Gi/o GPCRs. To determine if the promiscuous G α 15 protein is coupling your Gi/o GPCR of interest to the NFAT-*bla* reporter construct, it is necessary for the Response Ratio of the cell line containing the GPCR to be statistically different (*e.g.*, using a Student's T-test) than that of the parental cell line. The Response Ratio is defined as the Blue/Green ratio of maximally stimulated cells divided by the Blue/Green ratio of unstimulated cells (see **Section 10, Data Analysis**). A Response Ratio of ≥ 1.5 for the transiently transfected cell line generally indicates that it is a suitable candidate for the creation of a stable cell line.

The transfection procedure uses Lipofectamine[™] 2000, available separately from Invitrogen (see **Section 4**). The substrate loading procedure uses LiveBLAzer[™]-FRET B/G Substrate Mixture (CCF4-AM), also available separately (see **Section 4**). If you are using alternative substrates (*e.g.*, ToxBLAzer[™] DualScreen or LyticBLAzer[™] Loading kits), follow the loading protocol provided with the product.

7.1 Quick Assay Reference Guides

For a more detailed protocol, see **Section 7.2**.

| | Unstimulated Wells | Stimulated Wells | Cell-free Wells |
|---|---|---|-------------------------------------|
| Step 1 Plate cells | 90 μ l cells in Growth Medium (25,000 cells/well) | 90 μ l cells in Growth Medium (25,000 cells/well) | 90 μ l Growth Medium (no cells) |
| Step 2 Incubate cells | Incubate in a humidified 37°C/5% CO ₂ incubator for 16–24 hours | | |
| Step 3 Transfect cells | Aspirate Growth Medium and replace with 75 μ l Transfection Mix. | | |
| Step 4 Incubate cells | Incubate in a humidified 37°C/5% CO ₂ incubator for 16–20 hours | | |
| Step 5 Remove Transfection Mix | Aspirate Transfection Mix and replace with 90 μ l Growth Media. | | |
| Step 6 Incubate cells | Incubate in a humidified 37°C/5% CO ₂ incubator for 16–24 hours | | |
| Step 7 Add Agonist or Test Compounds | 10 μ l Assay Medium | 10 μ l 10X Agonist in Assay Medium | 10 μ l Assay Medium |
| Step 8 Incubate cells | Incubate in a humidified 37°C/5% CO ₂ incubator for 4–6 hours | | |
| Step 9 Prepare 6X Substrate Mix | 6 μ l of 1 mM LiveBLAzer [™] -FRET B/G (CCF4-AM) Substrate + 60 μ l of Solution B, mix. Add 904 μ l of Solution C, mix. Add 30 μ l of Solution D, mix. | | |
| Step 10 Add Substrate Mixture | 20 μ l per well | | |
| Step 11 Incubate Substrate Mix. + cells | 2 hours at room temperature in the dark | | |
| Step 12 Detect activity | See Section 9 | | |
| Step 13 Analyze data | See Section 10 | | |

For Technical Support for this or other Drug Discovery Products, dial 760-603-7200, option 3, extension 40266

7.2 Detailed Assay Protocol

We suggest using a minimum of 15 wells for the Transfected/Stimulated condition, the Transfected/Unstimulated condition, and the Cell-free Control.

7.2.1 Precautions

- Work on a dust-free, clean surface. Always handle the 96-well, black-wall, clear-bottom assay plate by the sides; do not touch the clear bottom of the assay plate.
- If pipetting manually, you may need to centrifuge the plate briefly at room temperature (for 1 minute at $14 \times g$) after additions to ensure all assay components are on the bottom of the wells.

7.2.2 Plating Cells

Day 1 (Plate the cells the day before the transfection):

1. Harvest cells and resuspend in Growth Medium (+) at a density of 2.7×10^5 cells/ml.
2. Add 90 μ l per well of the Growth Medium (+) to the Cell-free Control wells. Add 90 μ l per well of the cell suspension to the Stimulated wells and Unstimulated Control wells.

Note: Plate cells the day before the transfection and place them in a humidified 37°C/5% CO₂ incubator overnight.

7.2.3 Transient Transfection Procedure

Days 2 and 3:

Use the following procedure to transfect your Gi/o GPCR of interest into mammalian cells in a 96-well format. This procedure will make enough Transfection Mix for 33 wells.

Note: The plasmid containing your Gi/o GPCR of interest should *not* contain a Zeocin™ or Blasticidin resistance gene. We recommend using pcDNA™ 6.2-hygro-DEST Vector (see **Section 4** for ordering information).

1. **For each transfection sample**, prepare complexes as follows:
 - a. Dilute 4 μ g of the Gi/o coupled GPCR expression plasmid DNA in 250 μ l of Opti-MEM® I Reduced Serum Medium (or other medium without serum). Mix gently.
 - b. Mix Lipofectamine™ 2000 gently before use, then dilute 10 μ l in 250 μ l of Opti-MEM® I Medium. Incubate for 5 minutes at room temperature.

Note: Proceed to Step C within 25 minutes.

- c. After the 5 minute incubation, combine the diluted DNA with diluted Lipofectamine™ 2000 (total volume = 500 μ l). Mix gently and incubate for 20 minutes at room temperature (solution may appear cloudy).

Note: Complexes are stable for 6 hours at room temperature.

2. Add the 500 μ l of DNA-Lipofectamine™ 2000 complexes to 2 ml of Growth Medium (-). This is the transfection mix.
3. Remove Growth Medium from the wells of the plate and replace with 75 μ l of Transfection Mix per well. Incubate the assay plate in a humidified 37°C/5% CO₂ incubator for 16–20 hours.
4. Replace the Transfection Mix after 16–20 hours with 90 μ l of Growth Medium (+) per well. Incubate the assay plate in a humidified 37°C/5% CO₂ incubator for another 16–24 hours for high-level expression of the GPCR.

7.2.4 Agonist Test Assay Plate Setup

Day 4:

Following transfection, prepare an assay using a known agonist to test for a positive beta-lactamase response from the NFAT response element.

1. Prepare a 10X stock of a known agonist in Assay Medium.
2. Add 10 μ l of the 10X stock of agonist to the Stimulated wells.
3. Add 10 μ l of the Assay Medium alone to the Unstimulated Control wells and the Cell-free Control wells.
4. Incubate the agonist assay plate in a humidified 37°C/5% CO₂ incubator for 4–6 hours.

7.2.5 Substrate Loading

The following procedure uses LiveBLAzer[™]-FRET B/G Substrate Mixture (CCF4-AM), available separately from Invitrogen (see **Section 4**). If you are using alternative substrates (*e.g.*, ToxBLAzer[™] DualScreen or LyticBLAzer[™] Loading kits), follow the loading protocol provided with the product.

Prepare the LiveBLAzer[™]-FRET B/G Substrate Mixture (CCF4-AM) and load the cells in the absence of direct strong lighting. Turn off the light in the hood.

1. Prepare Solution A: 1 mM LiveBLAzer[™]-FRET B/G Substrate (CCF4-AM) Substrate Mixture in dry DMSO by adding 912 μ l of DMSO per mg of dry substrate. Store the aliquots of the stock solution at -20°C until use. The molecular weight of the LiveBLAzer[™]-FRET B/G Substrate (CCF4-AM) is 1096 g/mol.
2. Prepare 6X Loading Solution:
 - a. Add 6 μ l of Solution A to 60 μ l of Solution B and vortex.
 - b. Add 904 μ l of Solution C to the above solution and vortex.
 - c. Add 30 μ l of Solution D to the above solution and vortex.
3. Remove assay plate from the humidified $37^{\circ}\text{C}/5\% \text{CO}_2$ incubator.

Note: Handle the plate gently and do not touch the bottom.
4. Add 20 μ l (transient transfection assay plate) or 8 μ l (parental cell negative control assay plate) of the 6X Substrate Mixture to each well.
5. Cover the plate to protect it from light and evaporation.
6. Incubate at room temperature for 2 hours.
7. Proceed to **Section 9, Detection**.

7.3 Creating a Stable Cell Line

A stable pool of transfected cells can be created by selection with hygromycin. We recommend using a concentration of 100 $\mu\text{g}/\text{ml}$ of hygromycin for selection. Upon creation of a stable pool of cells containing the GPCR of interest, we recommend that you utilize flow cytometry to rapidly and quantitatively isolate a single clone with the desired performance characteristics (*e.g.*, maximum response ratio or moderate constitutive activity) for your experimental requirements.

8. Parental Cell Negative Control Assay Procedure

Use the following procedure to determine the activity of agonist compounds on the parental cell line. The following procedure uses LiveBLAzer[™]-FRET B/G Substrate Mixture (CCF4-AM), available separately from Invitrogen (see **Section 4**). If you are using alternative substrates (*e.g.*, ToxBLAzer[™] DualScreen or LyticBLAzer[™] Loading kits), follow the loading protocol provided with the product.

8.1 Quick Assay Reference Guides

For a more detailed assay protocol, see **Section 8.2**.

| | Unstimulated Wells | Stimulated Wells | Cell-free Wells | Test Compound Wells |
|--|---|---|---------------------------------------|---|
| Step 1 Plate cells, incubate | 32 μ l cells in Assay Medium (5,000 cells/well) | 32 μ l cells in Assay Medium (5,000 cells/well) | 32 μ l Assay Medium (no cells) | 32 μ l cells in Assay Medium 5,000 cells/well |
| Incubate cells for 16–24 hrs. at 37°C/ 5%CO ₂ | | | | |
| Step 2 Add Agonist or Test Compounds | 8 μ l Assay Medium with 0.5% DMSO | 8 μ l 5X agonist in Assay Medium with 0.5% DMSO | 8 μ l Assay Medium with 0.5% DMSO | 8 μ l 5X Test Compounds in 0.5% DMSO |
| Step 3 Incubate cells | Incubate in a humidified 37°C/5% CO ₂ incubator for 5 hours | | | |
| Step 4 Prepare 6X Substrate Mix | 6 μ l of 1 mM LiveBLAzer [™] -FRET B/G (CCF4-AM) Substrate + 60 μ l of solution B, mix. Add 904 μ l of Solution C, mix. Add 30 μ l of Solution D, mix. | | | |
| Step 5 Add Substrate Mixture | 8 μ l per well | | | |
| Step 6 Incubate Substrate Mix. + cells | 2 hours at room temperature in the dark | | | |
| Step 7 Detect activity | See Section 9 | | | |
| Step 8 Analyze data | See Section 10 | | | |

8.2 Detailed Assay Protocol

Plate layouts and experimental outlines will vary; in screening mode, we recommend using at least three wells for each control: Unstimulated Control, Stimulated Control, and Cell-free Control.

Note: Some solvents may affect assay performance. Assess the effects of solvent before screening. The cell stimulation procedure described below is carried out in the presence of 0.1% DMSO to simulate the effect that a Test Compound's solvent might have on the assay. If you use other solvents and/or solvent concentrations, optimize the following assay conditions appropriately.

8.2.1 Precautions

- Work on a dust-free, clean surface. Always handle the 384-well, black-wall, clear-bottom assay plate by the sides; do not touch the clear bottom of the assay plate.
- If pipetting manually, you may need to centrifuge the plate briefly at room temperature (for 1 minute at 14 \times g) after additions to ensure all assay components are on the bottom of the wells.

8.2.2 Plating Cells

1. Harvest cells and resuspend in Assay Medium to a density of 156,250 cells/ml.
2. Add 32 μ l per well of the Assay Medium to the Cell-free Control wells. Add 32 μ l per well of the cell suspension to the Test Compound wells, the Unstimulated Control wells, and Stimulated Control wells. Incubate cells at 37°C/5% CO₂ for 16–24 hours.

8.2.3 Assay Plate Setup

1. Prepare a stock solution of 0.5% DMSO in Assay Medium.
2. Prepare a 5X stock of Test Compounds in Assay Medium with 0.5% DMSO.
3. Prepare a 5X stock of agonist in Assay Medium with 0.5% DMSO. We recommend running a dose response curve to determine the optimal concentration of the agonist solution.
4. Add 8 μ l of the stock solution of 0.5% DMSO in Assay Medium to the Unstimulated Control and Cell-free Control wells.

For Technical Support for this or other Drug Discovery Products, dial 760-603-7200, option 3, extension 40266

- Add 8 μ l of the 5X stock solution of agonist to the Stimulated Control wells.
- Add 8 μ l of the 5X stock of Test Compounds to the Test Compound wells.
- Incubate the Agonist assay plate in a humidified 37°C/5% CO₂ incubator for 5 hours.

8.2.4 Substrate Loading

The following procedure uses LiveBLAzer[™]-FRET B/G Substrate Mixture (CCF4-AM), available separately from Invitrogen (see **Section 4**). If you are using alternative substrates (*e.g.*, ToxBLAzer[™] DualScreen or LyticBLAzer[™] Loading kits), follow the loading protocol provided with the product.

Prepare LiveBLAzer[™]-FRET B/G Substrate Mixture (CCF4-AM) Substrate Mixture and load cells in the absence of direct strong lighting. Turn off the light in the hood.

- Prepare Solution A: 1 mM LiveBLAzer[™]-FRET B/G Substrate (CCF4-AM) Substrate Mixture in dry DMSO by adding 912 μ l of DMSO per mg of dry substrate. Store the aliquots of the stock solution at -20°C until use. The molecular weight of the LiveBLAzer[™]-FRET B/G Substrate (CCF4-AM) is 1096 g/mol.
- Prepare 6X Loading Solution:
 - Add 6 μ l of Solution A to 60 μ l of Solution B and vortex.
 - Add 904 μ l of Solution C to the above solution and vortex.
 - Add 30 μ l of Solution D to the above solution and vortex.
- Remove assay plate from the humidified 37°C/5% CO₂ incubator.

Note: Handle the plate gently and do not touch the bottom.
- Add 20 μ l (transient transfection assay plate) or 8 μ l (parental cell negative control assay plate) of the 6X Substrate Mixture to each well.
- Cover the plate to protect it from light and evaporation.
- Incubate at room temperature for 2 hours.
- Proceed to **Section 9, Detection**.

9. Detection

Make measurements at room temperature from the bottom of the wells, preferably in black-wall, clear-bottom assay plates with low fluorescence background. Before reading the plate, remove dust from the bottom with compressed air.

9.2.1 Instrumentation, Filters, and Plates

- Fluorescence plate reader with bottom reading capabilities.
- Recommended filters for fluorescence plate reader:

| | |
|--------------------|-----------|
| Excitation filter: | 409/20 nm |
| Emission filter: | 460/40 nm |
| Emission filter: | 530/30 nm |

9.2.2 Reading an Assay Plate

- Set the fluorescence plate reader to bottom-read mode with optimal gain and 5 reads.
- Allow the lamp in the fluorescence plate reader to warm up for at least 10 minutes before making measurements.
- Use the following filter selections:

| | Scan 1 | Scan 2 |
|---------------------------|--|--|
| Purpose: | Measure fluorescence in the Blue channel | Measure FRET signal in the Green channel |
| Excitation filter: | 409/20 nm | 409/20 nm |
| Emission filter: | 460/40 nm | 530/30 nm |

10. Data Analysis

10.1 Background Subtraction and Ratio Calculation

We recommend that you subtract the background for both emission channels (460 nm and 530 nm).

1. Use the assay plate layout to identify the location of the Cell-free Control wells. These Control wells are used for background subtraction.
2. Determine the average emission from the Cell-free Control wells at both 460 nm (Average Blue Background) and 530 nm (Average Green Background).
3. Subtract the Average Blue background from all of the Blue emission data.
4. Subtract the Average Green background from all of the Green emission data.
5. Calculate the Blue/Green Emission Ratio for each well, by dividing the background-subtracted Blue emission values by the background-subtracted Green emission values.

10.2 Visual Observation of Intracellular Beta-lactamase Activity Using LiveBLAzer[™]-FRET B/G Substrate (CCF4-AM)

Note: Microscopic visualization of cells will cause photobleaching. Always read the assay plate in the fluorescence plate reader before performing microscopic visualization.

An inverted microscope equipped for epifluorescence and with either a xenon or mercury excitation lamp may be used to view the LiveBLAzer[™]-FRET B/G Substrate (CCF4-AM) signal in cells. To visually inspect the cells, you will need a long-pass filter passing blue and green fluorescence light, so that your eye can visually identify whether the cells are fluorescing green or blue.

Recommended filter sets for observing beta-lactamase activity are described below and are available from Chroma Technologies (800-824-7662, www.chroma.com).

Chroma Set # 41031

| | |
|--------------------|--------------------------|
| Excitation filter: | HQ405/20x (405 \pm 10) |
| Dichroic mirror: | 425 DCXR |
| Emission filter: | HQ435LP (435 long-pass) |

Filter sizes vary for specific microscopes and need to be specified when the filters are ordered. For epifluorescence microscopes, a long-pass dichroic mirror is needed to separate excitation and emission light and should be matched to the excitation filter (to maximally block the excitation light around 405 nm, yet allow good transmission of the emitted light).

10. References

1. Zlokarnik, G., *et al*, **Quantitation of Transcription and Clonal Selection of Single Living Cells with Beta-Lactamase as Reporter**, (1998) *Science*; **279**: p84-88.
2. Kunapuli P, Ransom R, Murphy K, Pettibone D, Kerby J, Grimwood S, Zuck P, Hodder P, Lacson R, Hoffman I, Inglese J, Strulovici B, **Development of an Intact Cell Reporter Gene Beta-lactamase Assay for G Protein-coupled Receptors**, (2003) *Analytical Biochem.*; **314**: p16-29.
3. Xing, H., Pollok, B., *et al*, **A Fluorescent Reporter Assay For The Detection of Ligands Acting Through G1 Protein-coupled Receptors**, (2000) *J. Receptor & Signal Transduction Research*; **20**: p189-210.
4. Qureshi, S., *et al*, **A One-Arm Homologous Recombination Approach for Developing Nuclear Receptor Assays in Somatic Cells**, (2003) *Assay and Drug Dev. Tech.*; **1**: p755-766.
5. Peekhaus, N. *et al*, **A Beta-Lactamase-Dependent Gal4-Estrogen Receptor Transactivation Assay for the Ultra-High Throughput Screening of Estrogen Receptor Agonists in a 3,456-Well Format**, (2003) *Assay and Drug Dev Tech*, **1**: p789-800.
6. Chin, J., *et al*, **Miniaturization of Cell-Based, Beta-Lactamase-Dependent FRET Assays to Ultra-High Throughput Formats to Identify Agonists of Human Liver X Receptors**, (2003) *Assay and Drug Dev. Tech.*; **1**: p777-787.
7. Whitney M, Rockenstein E, Cantin G, Knapp T, Zlokarnik G, Sanders P, Durick K, Craig FF, Negulescu PA., **A Genome-wide Functional Assay of Signal Transduction in Living Mammalian Cells**, (1998) *Nat. Biotechnol.*; **16**: p1329-1333.

11. Purchaser Notification

Limited Use Label License No. 5: Invitrogen Technology

The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. The buyer may transfer information or materials made through the use of this product to a scientific collaborator, provided that such transfer is not for any Commercial Purpose, and that such collaborator agrees in writing (a) not to transfer such materials to any third party, and (b) to use such transferred materials and/or information solely for research and not for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For products that are subject to multiple limited use label licenses, the terms of the most restrictive limited use label license shall control. Life Technologies Corporation will not assert a claim against the buyer of infringement of patents owned or controlled by Life Technologies Corporation which cover this product based upon the manufacture, use or sale of a therapeutic, clinical diagnostic, vaccine or prophylactic product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. If the purchaser is not willing to accept the limitations of this limited use statement, Life Technologies is willing to accept return of the product with a full refund. For information about purchasing a license to use this product or the technology embedded in it for any use other than for research use please contact Out Licensing, Life Technologies, 5791 Van Allen Way, Carlsbad, California 92008; Phone (760) 603-7200 or e-mail outlicensing@lifetech.com

Limited Use Label License No. 51: Blasticidin and the Blasticidin Selection Marker

Blasticidin and the blasticidin resistance gene (*bsd*) are the subject of U.S. Patent No. 5,527,701 sold under patent license for research purposes only. For information on purchasing a license to this product for purposes other than research, contact Licensing Department, Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, California 92008. Phone (760) 603-7200. Fax (760) 602-6500.

Limited Use Label License No. 108: Lentiviral Technology

The Lentiviral Technology (based upon the lentikat™ system) is licensed from Cell Genesys, Inc., under U.S. Patent Nos. 5,834,256; 5,858,740; 5,994,136; 6,013,516; 6,051,427; 6,165,782 and 6,218,187 and corresponding patents and applications in other countries for internal research purposes only. Use of this technology for gene therapy applications or bioprocessing other than for non-human research use requires a license from Cell Genesys (Cell Genesys, Inc. 342 Lakeside Drive, Foster City, California 94404). The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity), including non-gene therapy research and target validation applications in laboratory animals.

Limited Use Label License No. 150: GeneBLAzer® Technology

This product and/or its use is the subject of one or more of U.S. Patent Nos. 5,741,657, 5,955,604, 6,291,162, and 6,472,205 and foreign equivalents licensed to Life Technologies Corporation. The right to use this product for internal research specifically excludes the right to use this product to identify, discover, and profile compounds that act as a flavor, fragrance or taste-enhancers and modify a target identified in taste, olfaction, or pheromone detection, which compound does not require FDA approval of an NDA for claims of safety and efficacy. The right to use methods claimed in the foregoing patents with this product for research purposes can only be acquired by the use of GeneBLAzer® substrates purchased from Life Technologies Corporation or its authorized distributors.

Limited Use Label License No. 238: NFAT Technology

This product is the subject of one or more of US Patents 5,837,840, 6,096,515, 6,197,925, 6,150,099, pending applications and foreign equivalents licensed by Life Technologies Corporation.

Limited Use Label License No. 317: LentiVector® Technology

This product is licensed under U.S. Pat. Nos. 5,817,491; 5,591,624; 5,716,832; 6,312,682; 6,669,936; 6,235,522; 6,924,123 and foreign equivalents from Oxford BioMedica (UK) Ltd., Oxford, UK, and is provided for use in academic and commercial in vitro and in vivo research for elucidating gene function, and for validating potential gene products and pathways for drug discovery and development, but excludes any use of LentiVector® technology for: creating transgenic birds for the purpose of producing useful or valuable proteins in the eggs of such transgenic birds, the delivery of gene therapies, and for commercial production of therapeutic, diagnostic or other commercial products not intended for research use where such products do not consist of or incorporate a lentiviral vector. Information about licenses for commercial uses excluded under this license is available from Oxford BioMedica (UK), Ltd., Medawar Centre, Oxford Science Park, Oxford OX4 4GA UK enquiries@oxfordbiomedica.co.uk or BioMedica Inc 11622 EI Camino Real #100, San Diego CA 92130- 2049 USA. LentiVector is a registered US and European Community trade mark of Oxford BioMedica plc.

Use of Genetically Modified Organisms (GMO)

Information for European Customers The $G\alpha 15$ -NFAT-*bla* CHO-K1 cell lines are genetically modified to contain pLentibsd/ $G\alpha 15$ and p3XNFATBlaX-Zeo. As a condition of sale, use of this product must be in accordance with all applicable local legislation and guidelines including EC Directive 90/219/EEC on the contained use of genetically modified organisms.

Zeocin™ is a trademark of CAYLA Corporation.

© 2008, 2010 Invitrogen Corporation. All rights reserved. Reproduction forbidden without permission.