TaqMan® RNase P Assays/TaqMan® GAPDH Assays

QSY™ Quenched

Catalog Numbers 4485712, 4485713, 4485714, 4485715, A30064, and A30065

Pub. No. MAN0010239 Rev. B.0

Note: For safety and biohazard guidelines, see the "Safety" appendix in the *TaqMan* Multiplex PCR Optimization User Guide (Pub. No. MAN0010189). Read the Safety Data Sheets (SDSs) and follow the handling instructions. Wear appropriate protective eyewear, clothing, and gloves.

Product description

TaqMan® RNase P Assays and TaqMan® GAPDH Assays are optimized to detect expression levels of RNase P and GAPDH, respectively. These reference assays are optimized for the human genome. They use $QSY^{\mathbb{N}}$ as a quencher and $ABY^{\mathbb{N}}$, $JUN^{\mathbb{N}}$ or $VIC^{\mathbb{N}}$ as the 5' dye.

Note: JUN^{$^{\text{IM}}$} and ROX^{$^{\text{IM}}$} dyes cannot be used together because their emission spectra overlap.

See TaqMan® Multiplex PCR Optimization User Guide (Pub. No. MAN0010189) for more information.

TaqMan® Multiplex Master Mix (Cat. No. 4461881) is recommended.

Contents and storage

All assays provide 250 reactions in a 20 μL reaction volume.

Thaw reagents completely before use.

Assay	Cat. No.	Contents	Storage
TaqMan® RNase P Assay, VIC™ dye/QSY™ probe, 20X	A30064	Two unlabeled primers at 18 uM (900 nM final concentration) each QSY™ probe (VIC™ dye labeled) at 5 uM (250 nM final concentration)	-25°C to -15°C
TaqMan® RNase P Assay, VIC™ dye/QSY™ probe, primer-limited, 20X	A30065	Two unlabeled primers at 3 uM (150 nM final concentration) each QSY™ probe (VIC™ dye labeled) at 3 uM (150 nM final concentration)	
TaqMan® RNase P Assay, ABY™ dye/QSY™ probe, 20X	4485714	Two unlabeled primers at 18 uM (900 nM final concentration) each QSY™ probe (ABY™ dye labeled) at 5 uM (250 nM final concentration)	
TaqMan® RNase P Assay, ABY™ dye/QSY™ probe, primer-limited, 20X	4485715	Two unlabeled primers at 3 uM (150 nM final concentration) each QSY™ probe (ABY™ dye labeled) at 3 uM (150 nM final concentration)	Minimize freeze/thaw cycles.
TaqMan® GAPDH Assay, JUN™ dye/QSY™ probe, 20X	4485712	Two unlabeled primers at 18 uM (900 nM final concentration) each QSY™ probe (JUN™ dye labeled) at 5 uM (250 nM final concentration)	
TaqMan® GAPDH Assay, JUN™ dye/QSY™ probe, primer-limited, 20X	4485713	Two unlabeled primers at 3 uM (150 nM final concentration) each QSY™ probe (JUN™ dye labeled) at 3 uM (150 nM final concentration)	





The information in this guide is subject to change without notice.

DISCLAIMER: TO THE EXTENT ALLOWED BY LAW, THERMO FISHER SCIENTIFIC INC. AND/OR ITS AFFILIATE(S) WILL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, INDIRECT, PUNITIVE, MULTIPLE, OR CONSEQUENTIAL DAMAGES IN CONNECTION WITH OR ARISING FROM THIS DOCUMENT, INCLUDING YOUR USE OF IT.

Revision history: Revision history of Pub. No. MAN0010239

Revision	Date	Description
B.0	4 May 2018	Updated storage temperatures to match product labels.
		Included final concentrations of primers and probes.
		Added assays.
		Updated for general style, formatting, and branding.
A.0	7 April 2014	New document

Important Licensing Information: These products may be covered by one or more Limited Use Label Licenses. By use of these products, you accept the terms and conditions of all applicable Limited Use Label Licenses.

©2018 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific and its subsidiaries unless otherwise specified. TaqMan is a registered trademark of Roche Molecular Systems, Inc., used under permission and license.

