

# CD71 (Transferrin Receptor) Monoclonal Antibody (OKT9 (OKT-9)), eBioscience™

Product Details	
Size	100 µg
Species Reactivity	Human
Published Species	Human, Mouse
Host/Isotype	Mouse / IgG1, kappa
Class	Monoclonal
Type	Antibody
Clone	OKT9 (OKT-9)
Conjugate	Unconjugated
Form	Liquid
Concentration	0.5 mg/mL
Purification	Affinity chromatography
Storage buffer	PBS, pH 7.2
Contains	0.09% sodium azide
Storage conditions	4° C
RRID	AB_467338

Applications	Tested Dilution	Publications
Western Blot (WB)	1:1,000	-
Immunohistochemistry (IHC)	-	1 Publication
Immunocytochemistry (ICC/IF)	5 µg/mL	2 Publications
Flow Cytometry (Flow)	0.5 µg/test	13 Publications
Functional Assay (FN)	-	1 Publication

## Product Specific Information

Description: The OKT9 monoclonal antibody reacts with human CD71, a 170-180 kDa type II transmembrane protein. CD71, the transferrin receptor, exists as a homodimer on the cell surface and is essential for cellular growth. CD71 is expressed by immature proliferating cells and at low levels on resting mature lymphocytes. Antigen or mitogen stimulation of T and B cells upregulates the expression of CD71.

Applications Reported: The OKT9 (OKT-9) antibody has been reported for use in flow cytometric analysis.


Applications Tested: This OKT9 (OKT-9) antibody has been tested by flow cytometric analysis of unstimulated and CD3/CD28-stimulated (3 days) human blood cells. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Purity: Greater than 90%, as determined by SDS-PAGE.

Aggregation: Less than 10%, as determined by HPLC.

Filtration: 0.2 µm post-manufacturing filtered.


## Advanced Verification Data



### CD71 (Transferrin Receptor) Antibody (14-0719-82)


Altered expression of target protein upon cell treatment demonstrates antibody specificity. Immunofluorescence analysis of CD71 using Anti-CD71 Mouse Monoclonal Antibody (OKT9 (OKT-9)) (Product # 14-0719-80) shows increased expression of proteins in THP-1 cell line upon differentiation into Macrophage (M0). Cell treatment validation info.

## Product Images For CD71 (Transferrin Receptor) Monoclonal Antibody (OKT9 (OKT-9)), eBioscience™



### CD71 (Transferrin Receptor) Antibody (14-0719-82) in ICC/IF

Immunofluorescence analysis of CD71 was performed using 70% confluent log phase THP-1 cells differentiated into Macrophage (M0). The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 15 minutes, and blocked with 1% BSA for 1 hour at room temperature. The cells were labeled with CD71(Transferrin Receptor) Mouse Monoclonal Antibody (OKT9 (OKT-9)) (Product # 14-0719-80) at 5 µg/mL in 0.1% BSA, incubated at 4 degree Celsius overnight and then labeled with Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate (Product # A28175) at a dilution of 1:2000 for 45 minutes at room temperature (Panel a: green). Nuclei (Panel b: blue) were stained with ProLong™ Diamond Antifade Mountant with DAPI (Product # P36962). F-actin (Panel c: red) was stained with Rhodamine Phalloidin (Product # R415, 1:300). Panel d represents the merged image showing membrane localization. Panel e shows untreated cells with no signal. Panel f represents control cells with no primary antibody to assess background. The images were captured at 60X magnification.



### CD71 (Transferrin Receptor) Antibody (14-0719-82) in WB

Knockdown of CD71 was achieved by transfecting HeLa cells with SCRIB specific siRNAs (Silencer® select Product # s725\*). Western blot analysis (Fig. a) was performed using membrane enriched cell extracts from the CD71 knockdown cells (lane 3), non-specific scrambled siRNA transfected cells (lane 2) and untransfected cells (lane 1). The blot was probed with CD71 (Transferrin Receptor) Monoclonal Antibody OKT9 (OKT-9) (Product # 14-0719-80, 1:1000 dilution) and Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, HRP conjugate (Product # A28177, 0.25 µg/mL, 1:4000 dilution). Densitometric analysis of this western blot is shown in histogram (Fig. b). Decrease in signal upon siRNA mediated knock down confirms that antibody is specific to CD71.

View more figures on [thermofisher.com](http://thermofisher.com)

## 17 References

### Immunohistochemistry (1)

PloS one

#### Ferristatin II promotes degradation of transferrin receptor-1 in vitro and in vivo.

"14-0719 was used in Immunohistochemistry-immunofluorescence to show that ferristatin II degrades transferrin receptor-1 through a nystatin-sensitive lipid raft pathway in rats."

Authors: Byrne SL,Buckett PD,Kim J,Luo F,Sanford J,Chen J,Enns C,Wessling-Resnick M

**Species**  
Human

**Dilution**  
Not Cited

**Year**  
2014

### Immunocytochemistry (2)

Oncogene

#### Terminal differentiation and loss of tumorigenicity of human cancers via pluripotency-based reprogramming.

"Published figure using CD71 (Transferrin Receptor) monoclonal antibody (Product # 14-0719-82) in Immunofluorescence"

Authors: Zhang X,Cruz FD,Terry M,Remotti F,Matushansky I

**Species**  
Not Applicable

**Dilution**  
Not Cited

**Year**  
2013

Glycobiology

#### Epitope mapping, expression and post-translational modifications of two isoforms of CD33 (CD33M and CD33m) on lymphoid and myeloid human cells.

"14-0719 was used in Immunofluorescence to determine the expression and localisation of the two CD33 isoforms on several hematopoietic cell lines."

Authors: Pérez-Oliva AB,Martínez-Esparza M,Vicente-Fernández JJ,Corral-San Miguel R,García-Peñarrubia P,Hernández-Caselles T

**Species**  
Human

**Dilution**  
Not Cited

**Year**  
2011

### Flow Cytometry (13)

eLife

#### Gene-centric functional dissection of human genetic variation uncovers regulators of hematopoiesis.

"Published figure using CD71 (Transferrin Receptor) monoclonal antibody (Product # 14-0719-82) in Flow Cytometry"

Authors: Nandakumar SK,McFarland SK,Mateyka LM,Lareau CA,Ulirsch JC,Ludwig LS,Agarwal G,Engreitz JM,Przychodzen B,McConkey M,Cowley GS,Doench JG,Maciejewski JP,Ebert BL,Root DE,Sankaran VG

**Species**  
Not Applicable

**Dilution**  
Not Cited

**Year**  
2019

[View more Flow references on thermofisher.com](#)

### More applications with references on thermofisher.com

## FN (1)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.