invitrogen



# **Nestin Monoclonal Antibody (10C2)**


| <b>Product Details</b> |  |
|------------------------|--|
| Size                   | 100 μg   |
| Species Reactivity     | Human, Mouse   |
| Published Species      | Human, Mouse   |
| Host/Isotype           | Mouse / IgG1, kappa                                      |
| Class                  | Monoclonal   |
| Туре                   | Antibody   |
| Clone                  | 10C2   |
| Conjugate              | Unconjugated   |
| Immunogen              | Human nestin protein corresponding to residues 1464-1614 |
| Form                   | Liquid   |
| Concentration          | 1 mg/mL  |
| Purification           | Protein A  |
| Storage buffer         | PBS with 1mg/mL BSA, 30% glycerol                        |
| Contains               | 0.05% sodium azide                                       |
| Storage conditions     | -20°C  |
| RRID                   | AB_2536821   |

| Applications                              | Tested Dilution | Publications   |
|---|-----------------|----------------|
| Western Blot (WB)                         | 1:500-1:1,500   | 1 Publication  |
| Immunohistochemistry (IHC)                | -               | 1 Publication  |
| Immunohistochemistry (Paraffin) (IHC (P)) | 1:20-1:100      | 2 Publications |
| Immunohistochemistry (Frozen) (IHC (F))   | Assay-dependent | -              |
| Immunocytochemistry (ICC/IF)              | 1:100           | 9 Publications |
| Flow Cytometry (Flow)                     | 1 μg/test       | -              |

## **Product Specific Information**

MA1-110 has been successfully utilized in Western Blot, immunocytochemistry, flow cytometry, and immunofluorescence applications in human samples. MA1-110 also acts a neuronal stem cell marker and detects a band in Western blot at ~250 kDa.


## Advanced Verification Data



## **Nestin Antibody (MA1-110)**


The specificity of anti-Nestin monoclonal antibody (Product # MA1-110) was demonstrated in western blot analysis by the detection of endogenous Nestin protein expression in human neural stem cells, but not in negative control line HepG2 which is not known to express Nestin protein. Relative expression validation info.

## **Product Images For Nestin Monoclonal Antibody (10C2)**



#### Nestin Antibody (MA1-110) in ICC/IF

Immunofluorescent analysis of Nestin (green) in human neural stem cells derived from PD-3 iPSCs using Gibco® PSC Neural Induction Medium (Product # A1647801). The cells were fixed and permeabilized using Image-IT® Fixation /Permeabilization kit (Product # R37602), and blocked with the including blocking buffer for one hour at room temperature. Cells were stained with a Nestin monoclonal antibody (Product # MA1-110) at a dilution of 1:100 in blocking buffer for 3 hours at room temperature, and then incubated with a DyLight488-conjugated goat antimouse IgG secondary antibody (Product # 35502) at a dilution of 1:250 for 1 hour at room temperature. Nucleus DNA (blue) was stained with DAPI (Product # D1306). Images were taken on an EVOS® FLoid® Cell Imaging Station at 10X magnification.



#### Nestin Antibody (MA1-110) in IHC

Immunohistochemistry was performed on human kidney tissue. To expose target protein, antigen was retreived using 10mM sodium citrate followed by microwave treatment for 8-15 minutes. Endogenous peroxidases were blocked in 3% H202-methanol for 15 minutes and tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with a nestin Rabbit polyclonal antibody (Product # MA1-110) at a dilution of 1:100 overnight in a humidified chamber. Tissues were washed in PBST and detection was performed using a secondary antibody conjugated to HRP. DAB staining buffer was applied and tissues were counterstained with hematoxylin and prepped for mounting. Images were taken at 40X magnification.

View more figures on thermofisher.com

#### **□** 13 References

#### Western Blot (1)

Frontiers in cell and developmental biology

CPNE7-Induced Autophagy Restores the Physiological Function of Mature Odontoblasts.

"Published figure using Nestin monoclonal antibody (Product # MA1-110) in Western Blot"

Authors: Park YH, Son C, Seo YM, Lee YS, Har A, Park JC

**Species**Not Applicable

**Dilution** Not Cited

**Year** 2021

### Immunohistochemistry (1)

Histochemistry and cell biology

Androgen receptors and gender-specific distribution of alkaline phosphatase in human thyroid cartilage.

"MA1-110 was used in immunohistochemistry to study gender differences in the localization of android receptors and alkaline phosphatase in human thyroid cartilage"

Authors: Claassen H, Mönig H, Sel S, Werner JA, Paulsen F

Species Human

Dilution 1:20

**Year** 2006

## Immunohistochemistry (Paraffin) (2)

**Human pathology** 

Comparative immunomorphology of testicular Sertoli and sertoliform tumors.

"MA1110 was used in immunohistochemistry - paraffin section to develop methods to distinguish Sertoli cell tumors from sertoliform tumors"

Authors: Mesa H,Gilles S,Datta MW,Murugan P,Larson W,Dachel S,Manivel JC

Species Human

Dilution

1:300

**Year** 2017

PloS one

Embryonic stem cells markers SOX2, OCT4 and Nanog expression and their correlations with epithelial-mesenchymal transition in nasopharyngeal carcinoma.

"MA1-110 was used in immunohistochemistry - paraffin section to assess the expression and prognostic impact of embryonic stem cell markers in nasopharyngeal carcinoma patients"

Authors: Luo W,Li S,Peng B,Ye Y,Deng X,Yao K

**Species** Not Applicable

Dilution 1:25

**Year** 2013

#### More applications with references on thermofisher.com

## ICC/IF (9)

For Research Use Only, Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation,"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended uses not restend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely likestrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. No O'THER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES, OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERFOLO IS LIMITED TO REPEAR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS) AT SHEE IS ON EXPRESS OR OR OR SHEED THE ORD FOR ANY SHEET OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses, or any type of consumption to human or animals.